

Interaction effects on magnetooscillations in a two-dimensional electron gas

Y. Adamov^{1,2}, I. V. Gornyi^{2†}, and A. D. Mirlin^{2,3#}

¹ *Department of Physics, Brookhaven National Laboratory, Upton, NY 11973-5000, USA*

² *Institut für Nanotechnologie, Forschungszentrum Karlsruhe, 76021 Karlsruhe, Germany*

³ *Institut für Theorie der Kondensierten Materie,
Universität Karlsruhe, 76128 Karlsruhe, Germany*

(Dated: February 2, 2008)

Motivated by recent experiments, we study the interaction corrections to the damping of magnetooscillations in a two-dimensional electron gas (2DEG). We identify leading contributions to the interaction-induced damping which are induced by corrections to the effective mass and quantum scattering time. The damping factor is calculated for Coulomb and short-range interaction in the whole range of temperatures, from the ballistic to the diffusive regime. It is shown that the dominant effect is that of the renormalization of the effective electron mass due to the interplay of the interaction and impurity scattering. The results are relevant to the analysis of experiments on magnetooscillations (in particular, for extracting the value of the effective mass) and are expected to be useful for understanding the physics of a high-mobility 2DEG near the apparent metal-insulator transition.

I. INTRODUCTION

The influence of electron-electron interaction on transport properties of low-dimensional disordered conductors at sufficiently low temperatures T remains one of central topics of the condensed matter physics. In a seminal set of works (see the review¹), Altshuler and Aronov studied the effects of the interplay of interaction and disorder on conductivity and tunneling density of states in the diffusive regime characterized by the condition $2\pi T\tau \ll 1$, where τ is the transport mean free time (we set $k_B = \hbar = 1$). Their results were generalized within the framework of the renormalization group (RG) by Finkelstein². The last decade has witnessed a renewed increase of activity in this field, largely motivated by experiments on an apparent metal-insulator transition in 2D systems. This interest was triggered by experiments³ which showed a “metallic” behavior (decrease of resistivity with lowering T) in high-mobility Si structures. Later, qualitatively similar behavior was observed in a variety of high-mobility 2D systems, see Refs. 4,5,6,7 for reviews.

The metallic behavior has been attributed to the effects of the electron-electron interaction in the ballistic temperature range, $2\pi T\tau \gg 1$. These effects were originally considered in the framework of the temperature-dependent screening⁸. More recently, a systematic theory was developed, taking into account also exchange contributions and the effects of both parallel and transverse magnetic fields, and valid in the whole range of T from the diffusive to the ballistic regime^{9,10}. Another mechanism that can explain the metallic behavior of resistivity in an intermediate temperature range in the diffusive regime was studied within the RG framework in Ref. 11. It is applicable to systems with more than one valley, such as silicon MOSFETs.

Despite these successes of the theory, numerous experimental observations remain puzzling and wait for an explanation. In particular, it was found that the spin susceptibility, proportional to the product mg of the effective mass m and the g -factor, is strongly growing when the density approaches the value n_c corresponding to the apparent transition. This conclusion was drawn on the basis of several experimental methods, including the analysis of beating pattern of Shubnikov-de Haas oscillations^{12,13,14,15}, the study of magnetoresistance^{16,17} in the parallel field, and measurement of thermodynamic magnetization¹⁸; see recent reviews^{19,20}.

The enhancement of susceptibility with lowering density, interpreted in a number of papers as its divergence at $n = n_c$, has attracted a great deal of attention, since it might be an indicator of some phase transition that the system undergoes with a decrease of density. The interpretation of the data has remained, however, controversial. In particular, it remained unclear whether the strong increase of spin susceptibility should be attributed to that of m or of g . This information is of crucial importance for understanding the nature of the possible transition.

Several experimental approaches have been used to separate the behavior of the effective mass from that of the g -factor. In Refs. 21,22,23 a fit of the resistivity data to the theoretical formulas of Ref. 9 was used to find the interaction constant F_0^σ , and thus the g -factor. The accuracy of this procedure is questionable, since the theory of Ref. 9 neglects higher Fermi-liquid interaction constants and assumes isotropic impurity scattering. Another approach is based on thermodynamic measurements in strong magnetic field²⁴. However, the authors of this work were not able to measure the effective mass in a very narrow interval of electron concentration only, so that the results are not too informative. Also, a strong magnetic field is expected to influence strongly the characteristics of the electron liquid, so that the applicability of such measurements to the low-field properties is questionable. So, while most of the above measurements seem to indicate that it is the effective mass that is responsible for the strongly enhanced susceptibility,

an independent verification is clearly needed.

A well known method for determination of the effective mass is based on the investigation of the temperature dependence of Shubnikov-de Haas oscillations (SdHO). It was applied to the present problem in Refs. 14,25. However, the analysis of the SdHO data is complicated by the fact that both the effective mass and the elastic quantum scattering time τ_q are T -dependent, in view of the combined effect of interaction and disorder. An unambiguous interpretation of experimental data requires¹⁴ a theoretical information on T -dependence of m and τ_q . A development of the corresponding theory is the aim of the present paper.

In fact, a recent paper²⁶ has made an important step in this direction. Specifically, it was shown in Ref. 26 that the Lifshitz-Kosevich formalism²⁷, originally developed for the analysis of magnetooscillations in a 3D Fermi liquid, is also applicable in 2D in the regime where the oscillations are exponentially suppressed by temperature smearing or disorder. (In the regime of strong oscillations, the Lifshitz-Kosevich formula in 2D should be modified, as was earlier shown in Ref. 28.) Another result of Ref. 26 is that the inelastic electron-electron relaxation does not contribute to the damping of magnetooscillations (similarly to the earlier result of Ref. 29 for the case of electron-phonon scattering).

The authors of Ref. 26 then calculated the contribution to the damping induced by the interplay of interaction and disorder. Their theoretical treatment of the problem is, however, far from complete. First, they consider only diagrams for the self-energy with one impurity-ladder vertex correction to the interaction line and discard diagrams with no and with two vertex corrections. Second, they claim that the T -dependence of the oscillation damping rate can be equivalently attributed either to the correction to the effective mass, or to the quantum scattering rate (Dingle temperature). Furthermore, in the latter case their result for the T -dependence of τ_q is in contradiction with the picture of Friedel oscillations inducing a correction to the relaxation rate, which is linear in T and is governed by backscattering⁹.

In addition to the above experimental motivation, the development of the theory of interaction effects on magnetooscillations in a disordered 2DEG represents a fundamental theoretical problem. Such a theory should complement the recently developed theory of interaction effects in transport of 2D electrons in zero and non-quantizing magnetic fields^{9,10}. Let us emphasize a peculiar aspect of the present problem. The damping of oscillations is governed by the self-energy, which is a single-particle quantity. [Indeed, the relevant diagrams, see Sec II B below, are reminiscent of those for the tunneling density of states (DOS)]. Generally, the self-energy is not a gauge-invariant object. On the other hand, the magnetization and the conductivity (magnetooscillations of which we would like to study) are observable (and thus gauge-invariant) quantities. It is well known that the gauge-invariance is of crucial importance for interaction-induced corrections; a difference between the results for conductivity and for tunneling DOS in the case of Coulomb interaction^{1,2} serves as a nice illustration. It is thus a theoretical challenge to see how the gauge invariance manifests itself in the magnetooscillation problem.

The outline of the article is as follows. The section II is devoted to presentation of the general formalism. In Sec. III we apply it to calculate the interaction-induced contribution to the damping of magnetooscillations in the case of short-range interaction. In Sec. IV we show how to extract from the above result the corrections to the effective mass and the quantum scattering time. We also perform a calculation of the correction to the scattering time based on the picture of Friedel oscillations and demonstrate a complete agreement between the two approaches. In Sec. V we generalize our results to the case of Coulomb interaction. We also perform there a comparison with the calculation of Ref. 26. Section VI summarizes our findings. Some technical details of our calculations are presented in Appendices.

II. MAGNETOOSCILLATIONS: GENERAL FORMALISM

A. Derivation of the formula for a decay of the oscillations

We begin by calculating the oscillatory part Ω_{osc} of the thermodynamic potential Ω . From this quantity one can derive the oscillating contribution to the thermodynamic density of states

$$\frac{\partial n_{\text{osc}}}{\partial \mu} = \frac{\partial^2 \Omega_{\text{osc}}}{\partial \mu^2}, \quad (1)$$

where μ is the chemical potential, and de Haas-van Alphen oscillations of magnetic susceptibility

$$\chi_{\text{osc}} = -\frac{\partial^2 \Omega_{\text{osc}}}{\partial B^2}, \quad (2)$$

where B is a magnetic field. The main subject of our interest is the exponential damping factor of these magnetooscillations. For non-interacting electrons, the same exponential damping factor governs the magnitude of the Shubnikov-de Haas oscillations of the conductivity for the case of weak disorder potential in sufficiently weak magnetic field^{35,36},

where the self-consistent Born approximation³⁵ (SCBA) is valid. As we are going to show, the interaction-induced correction to the damping factor of the thermodynamic density of states arises due to the renormalization of the effective mass and the quantum scattering time. Therefore these T -dependent corrections to the damping factor govern the magnitude of the Shubnikov – de-Hass oscillations as well, similarly to the non-interacting case.

Our starting point is the expression for the thermodynamic potential derived in the paper by Luttinger and Ward³⁰

$$\Omega = -T \text{Tr} \ln(-G^{-1}) - T \text{Tr}(G\Sigma) + \Omega'. \quad (3)$$

Here

$$G(i\varepsilon_n, m\omega_c) = [G_0^{-1}(i\varepsilon_n, m\omega_c) - \Sigma(i\varepsilon_n, m\omega_c)]^{-1}, \quad (4)$$

is the dressed Green's function in the Matsubara formalism, $i\varepsilon_n = (2n+1)i\pi T$ is the Matsubara fermionic energy, ω_c is the cyclotron frequency, and m is the Landau level index. Further,

$$G_0(i\varepsilon_n, m\omega_c) = \frac{1}{i\varepsilon_n + \mu - (m + 1/2)\omega_c}$$

is the Green's function in the absence of disorder and interaction, and $\Sigma(i\varepsilon_n, m\omega_c)$ is a self-energy part of Green's function which includes all the disorder and interaction effects.

The trace in Eq. (3) implies summation over Matsubara frequencies ε_n and over Landau levels m . The logarithmic term contains all the closed loop diagrams with insertion of self-energy (Fig. 1). The terms $-T \text{Tr}(G\Sigma)$ and Ω' are introduced to avoid double-counting of diagrams^{30,31}. The term Ω' denotes the sum of all so-called skeleton diagrams with all bare Green's functions replaced by dressed Green's functions (for the recent discussion of Luttinger-Ward formalism in 2D Fermi systems see Refs. 32,33).

As shown in Ref. 34, the exponential decay of magnetooscillations is described by the $\text{Tr} \ln$ -term. The oscillatory parts of the additional terms, which are introduced to fight overcounting, cancel each other. In order to obtain the correction to the thermodynamic potential we need to calculate the self-energy part of the Green's function.

We decompose the self-energy into two parts:

$$\Sigma(i\varepsilon_n, m\omega_c) = \Sigma_{\text{dis}}(i\varepsilon_n, m\omega_c) + \Sigma_{\text{ee}}(i\varepsilon_n, m\omega_c). \quad (5)$$

where $\Sigma_{\text{dis}}(i\varepsilon_n, m\omega_c)$ denotes the self-energy part due to the scattering on disorder potential with electron-electron interaction switched off and $\Sigma_{\text{ee}}(i\varepsilon_n, m\omega_c)$ contains all the interaction effects.

In this paper we assume that disorder potential is δ -correlated, inducing a large-angle scattering of electrons. The disorder-induced (noninteracting) part of the self-energy for white-noise disorder and weak magnetic field $\omega_c\tau \ll 1$ is given by

$$\Sigma_{\text{dis}}(i\varepsilon_n, m\omega_c) = \frac{i \text{sgn} \varepsilon_n}{2\tau}. \quad (6)$$

For stronger magnetic field (i.e. for separated Landau levels), one should employ the SCBA. In this case, both the real and imaginary parts of the self-energy depend on $i\varepsilon_n$ in a non-trivial way. In this paper, however, we will address only weak magnetic fields when Landau levels overlap.

$$-T \text{Tr} \ln(-G^{-1}) = -T \text{Tr} \ln(-G_0^{-1}) + \text{[diagram with 1 Sigma]} + \frac{1}{2} \text{[diagram with 2 Sigma]} + \frac{1}{3} \text{[diagram with 3 Sigma]} + \dots$$

FIG. 1: The logarithmic term $\tilde{\Omega}$, Eq. (7), in the thermodynamic potential, Eq. (3), is a sum of closed loop diagrams with self-energy insertions Σ . This term is responsible for magnetooscillations.

We thus consider the relevant term $\tilde{\Omega} = -T \text{Tr} \ln(-G^{-1})$ in the thermodynamic potential,

$$\tilde{\Omega} = -2\nu T \sum_{n=-\infty}^{\infty} \omega_c \sum_{m=0}^{\infty} \ln[\omega_c m - \mu - i\varepsilon_n + \Sigma(i\varepsilon_n, \omega_c m)]. \quad (7)$$

For overlapping Landau levels,

$$\omega_c \tau \ll 1, \quad (8)$$

the k -th harmonics of the magnetooscillations

$$\Omega_{\text{osc}} = \sum_k A_k \cos \frac{2k\pi^2 n_e}{eB}$$

is damped by disorder even at zero temperature via the standard Dingle factor $\exp(-\pi/\omega_c \tau)$. Therefore we will consider only the first harmonics of the oscillations, A_1 , neglecting all A_k with $k > 1$ (whose damping is much stronger).

The oscillatory part of (7) is calculated in Appendix A:

$$\Omega_{\text{osc}} \simeq 2\nu \left(\frac{\omega_c}{2\pi} \right)^2 A_1 \cos \frac{2\pi^2 n_e}{eB}, \quad (9)$$

(n_e is the electron concentration) with the amplitude of the first harmonics of the oscillatory part of the thermodynamic potential given by

$$A_1 \equiv \frac{4\pi^2 T}{\omega_c} \sum_{\varepsilon_n > 0} \exp \left(-\frac{2\pi}{\omega_c^*} \left[\varepsilon_n + \frac{1}{2\tau(1+\alpha_0)} + \frac{i\delta\Sigma(i\varepsilon_n, \xi_0)}{1+\alpha_0} \right] \right). \quad (10)$$

Here $\delta\Sigma(i\varepsilon_n, \xi_0)$ is the self-energy part related to the interplay of disorder and interaction. It is analytically continued from the points $m\omega_c$ to the whole complex plane ξ and taken at $\xi = \xi_0$, where ξ_0 [defined in Eqs. (A.9) and (A.10) of Appendix A] is the pole of the Green's function in the presence of disorder.

The coefficients β_0 and α_0 determine the Fermi-liquid (FL) renormalization of the effective mass in a pure system at zero T ,

$$m^* = m \frac{1+\alpha_0}{1+\beta_0}. \quad (11)$$

The effective mass m^* in turn governs the expression for the FL-renormalized effective cyclotron frequency,

$$\omega_c^* = \frac{eB}{m^*} = \omega_c \frac{1+\beta_0}{1+\alpha_0}. \quad (12)$$

The coefficient α_0 is related to the FL renormalization of the Z -factor,

$$Z = \frac{1}{1+\alpha_0}, \quad (13)$$

which is given by the residue of the Green's function.

Depending on the relation between temperature T and the elastic scattering rate $1/\tau$, there are two regimes: ballistic, $T\tau \gg 1$, and diffusive, $T\tau \ll 1$ (more accurately, the relevant dimensionless parameter is $2\pi T\tau$). In the ballistic regime, it follows from Eq. (8) that $T \gg 1/\tau \gg \omega_c$. The diffusive regime can be further split into two subregimes: normal diffusive ($\omega_c \ll T \ll 1/\tau$) and ultra-diffusive ($T \ll \omega_c \ll 1/\tau$). When $T \gg \omega_c$, as in the ballistic and normal diffusive regimes, only the first Matsubara frequency $\varepsilon_0 = \pi T$ in the sum determining A_1 is relevant, since the contribution of higher Matsubara frequencies are exponentially suppressed. On the other hand, in the ultra-diffusive regime $T \ll \omega_c$ and higher Matsubara frequencies contribute as well.

In what follows we concentrate on the case $T \gg \omega_c$. Under this condition, we get

$$A_1 = \frac{4\pi^2 T}{\omega_c} \exp \left[-\frac{2\pi}{\omega_c^*} \{ \pi T + i Z \delta\Sigma(i\pi T, \xi_0) \} \right] \exp \left[-\frac{\pi}{\omega_c^* \tau^*} \right], \quad (14)$$

where we introduced the FL-renormalized scattering time

$$\tau^* = \tau (1 + \alpha_0). \quad (15)$$

We note that this renormalization of τ is incomplete since it does not include the FL vertex corrections to the impurity scattering line. The corresponding contributions is contained in $\delta\Sigma(i\pi T, \xi_0)$ [T -independent terms in Eqs. (41) and




FIG. 2: Self energy diagrams in the first order in the effective interaction (wavy line). Black triangles denote impurity ladders Γ dressing interaction vertices, Fig. 3. (a) “Simple” self-energies Σ_{ij}^a ; (b) Hikami-box self-energies Σ_{ij}^b generated by covering each of Σ_{ij}^a by an impurity line (dashed).



FIG. 3: Interaction vertex renormalized by the impurity ladder, $\Gamma(i\omega_k, \mathbf{q})$. Dashed line represents scattering on impurity.

(81)] and will be addressed in Sections III and IV below. We also note that the inelastic contribution to the self-energy $\propto [(\pi T)^2 - \varepsilon_n^2] \text{sgn } \varepsilon_n$ (see Appendix B) vanishes for $\varepsilon_n = \pi T$, and thus does not affect the damping of the magnetooscillations $B(T)$ for $T \gg \omega_c$, in agreement with Refs. 26,29.

Using the renormalized quantities in Eq. (14) we represent A_1 in the form

$$A_1 = A_1^{(0)}(T) \exp[B(T)], \quad (16)$$

where

$$A_1^{(0)}(T) \equiv \frac{4\pi^2 T}{\omega_c} \exp \left[-\frac{2\pi^2 T}{\omega_c^*} - \frac{\pi}{\omega_c^* \tau^*} \right] \quad (17)$$

is the standard FL Lifshitz-Kosevich result and

$$B(T) = -\frac{2\pi i Z \delta \Sigma(i\pi T, \xi_0)}{\omega_c^*}. \quad (18)$$

We thus see that in order to obtain the additional interaction-induced damping factor of magnetooscillations it is necessary to evaluate $Z \delta \Sigma(i\pi T, \xi_0)$.

B. Self-energy

We begin by considering the interaction-induced self-energy part $\Sigma_{ee}(i\varepsilon_n, m\omega_c)$ in the lowest order in interaction. This is sufficient in the case of a weak short-range interaction analyzed in Sec. III below. For the more realistic case of the Coulomb interaction (Sec. V), the relevant higher-order terms can be treated using the random-phase approximation (RPA). Higher-order contributions to the T -dependent part of the self-energy, $\delta \Sigma(i\pi T, \xi_0)$, are small in the parameter $1/E_F \tau$ or T/E_F .

Let us list important elements which are necessary for calculation of the interaction-induced part of the self-energy (Fig. 2). Each contribution to the self-energy has exchange and Hartree parts. We first address the exchange contribution (the Hartree terms can be written in a similar way). It contains the angle-averaged Green's function covered by the effective interaction line. The corresponding vertices may be dressed by impurity ladders (Fig. 3). Notice that the renormalized vertex includes at least one impurity line. Finally, when the interaction line changes the signs of Matsubara frequencies at vertices, an additional diagram (we term it a “Hikami-box diagram”) with a single impurity line covering the whole block is to be included (Fig. 2b).

We split $\Sigma_{ee}(i\varepsilon_n, m\omega_c)$ into three contributions, corresponding to different possibilities of dressing the two interaction vertices by impurities,

$$\Sigma_{ee} = \Sigma_{00} + 2\Sigma_{01} + \Sigma_{11}. \quad (19)$$

Here the subscripts $i, j = 0, 1$ indicate whether the corresponding vertex is dressed by an impurity ladder. The factor 2 in front of Σ_{01} term reflects two possibilities of dressing one of the interaction vertices. Each of the terms Σ_{ij} is a sum of two contributions, $\Sigma_{ij} = \Sigma_{ij}^a + \Sigma_{ij}^b$, where Σ_{ij}^a is the “simple” self-energy (Fig. 2a) and Σ_{ij}^b is its Hikami-box counterpart (Fig. 2b). We note that in Ref. 26 only one out of six diagrams (namely, Σ_{01}^a) was taken into account.

The expression for Σ_{00}^a in a finite magnetic field reads

$$\Sigma_{00}^a(i\varepsilon_n, m\omega_c) = -T \sum_{\omega_k} \sum_L \int \frac{d^2q}{(2\pi)^2} J_{|L|}^2(qR_c) V(i\omega_k, \mathbf{q}) G(i\varepsilon_n - i\omega_k, m\omega_c + L\omega_c), \quad (20)$$

where

$$G(i\varepsilon_n, m\omega_c) = \frac{1}{i\varepsilon_n + \mu - (m + 1/2)\omega_c - \Sigma(i\varepsilon_n, m\omega_c)} \quad (21)$$

is the Green’s function in Landau levels representation, $V(i\omega_k, \mathbf{q})$ is the effective interaction, and $J_{|L|}^2(qR_c)$ describes the bare vertex function connecting Landau levels m and $m + L$ in the quasiclassical limit $m, m + L \gg 1$ (for details see, e.g., Ref. 37 and references therein). In the expressions for Σ_{01}^a and Σ_{11}^a this bare vertex function is multiplied by $\Gamma(i\omega_k, \mathbf{q})$ and $\Gamma^2(i\omega_k, \mathbf{q})$, respectively, where Γ is the impurity ladder (Fig.3).

In the limit $B \rightarrow 0$, the corresponding self-energies depend on the momentum $\mathbf{p}$ instead of the Landau level index m . The interaction vertices are dressed only when the Matsubara energies at the vertices have opposite signs, which restricts the summation over ω_k in Σ_{01} and Σ_{11} to the domain $\varepsilon_n(\omega_k - \varepsilon_n) > 0$:

$$\Sigma_{00}^a(i\varepsilon_n, \xi_p) = -T \sum_{\omega_k} \int \frac{d^2q}{(2\pi)^2} V(i\omega_k, \mathbf{q}) G(i\varepsilon_n - i\omega_k, \mathbf{p} - \mathbf{q}), \quad (22)$$

$$\Sigma_{01}^a(i\varepsilon_n, \xi_p) = -T \sum_{\varepsilon_n(\omega_k - \varepsilon_n) > 0} \int \frac{d^2q}{(2\pi)^2} V(i\omega_k, \mathbf{q}) \Gamma(i\omega_k, \mathbf{q}) G(i\varepsilon_n - i\omega_k, \mathbf{p} - \mathbf{q}), \quad (23)$$

$$\Sigma_{11}^a(i\varepsilon_n, \xi_p) = -T \sum_{\varepsilon_n(\omega_k - \varepsilon_n) > 0} \int \frac{d^2q}{(2\pi)^2} V(i\omega_k, \mathbf{q}) \Gamma^2(i\omega_k, \mathbf{q}) G(i\varepsilon_n - i\omega_k, \mathbf{p} - \mathbf{q}), \quad (24)$$

where $G(i\varepsilon_n, \mathbf{p}) = [i\varepsilon_n + \mu - \xi_p + i\text{sgn}\varepsilon_n/2\tau - \Sigma_{ee}(i\varepsilon_n, \xi_p)]^{-1}$ with $\xi_p = p^2/2m$ and the vertex correction (Fig.3) reads

$$\Gamma(i\omega_k, \mathbf{q}) = \frac{1}{\sqrt{(|\omega_k|\tau + 1)^2 + (qv_F\tau)^2} - 1}. \quad (25)$$

To calculate the damping factor of the oscillations, we need the self-energy taken at the value of ξ which is determined by the pole of the Green’s function, $\xi = \xi_0$, see Appendix B. According to Eq. (18), the self-energy is further multiplied by Z in the damping exponent. This is equivalent to calculating the following integral:

$$\begin{aligned} Z\Sigma_{ij}^a(i\varepsilon_n, \xi_0) &= \frac{\text{sgn}\varepsilon_n}{2\pi i} \frac{1 + \beta_0}{1 + \alpha_0} \int d\xi_k G(i\varepsilon_n, k) \Sigma_{ij}^a(i\varepsilon_n, \xi_k) \\ &\simeq \text{sgn}\varepsilon_n \frac{v_F^*}{2\pi i} \int dk G(i\varepsilon_n, k) \Sigma_{ij}^a(i\varepsilon_n, \xi_k). \end{aligned} \quad (26)$$

We note that the Z -factor drops out in the product $Z\Sigma$. Indeed, the Green’s function under the interaction line in the self-energy contains the Z -factor in the numerator so that in the numerator of the product $Z\Sigma$ we get the factor Z^2 . However, the Z -factor is not a gauge invariant quantity and therefore should not appear in the expressions for observables, in contrast to the FL-renormalized effective mass. At this point we should take into account the FL renormalization of the two interaction vertices in Σ . Since we are interested in the contribution of relatively slow transferred momenta and frequencies giving rise to the T dependence of $B(T)$, $q \ll k_F$ and $\omega_k \ll E_F$, we can set them to zero when considering the FL vertex renormalization. Then one can apply the Ward identity for the FL-interaction dressing of the vertices³⁹, which amounts to multiplying each vertex by a factor $1/Z$. These vertex factors

in the denominator of $B(T)$ cancel Z^2 in the numerator of $B(T)$. This implies that one can simply discard such renormalizations, setting $Z = 1$ everywhere, when the observable quantities are calculated. The FL renormalization then amounts to replacement of the bare band mass m , Fermi velocity v_F , and elastic scattering time, τ , by the renormalized parameters, m^* , v_F^* and τ^* , respectively. In what follows, we will omit the asterisks, using the notation m , v_F and τ for the renormalized quantities.

The relevant contributions to the self-energy are calculated in Appendix B. Combining all the terms together, we have

$$\delta\Sigma(i\varepsilon_n, \xi_0) = -i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2q}{(2\pi)^2} V(i\omega_m, \mathbf{q}) K(i\omega_m, \mathbf{q}), \quad (27)$$

$$K(i\omega_m, \mathbf{q}) = \frac{[1 + \Gamma(i\omega_m, \mathbf{q})]^2}{S(i\omega_m, \mathbf{q})} \left[1 - \frac{W}{\tau S^2(i\omega_m, \mathbf{q})} \right] - \frac{1}{S_0(i\omega_m, \mathbf{q})}, \quad (28)$$

where

$$S(i\omega_m, \mathbf{q}) = \sqrt{(|\omega_m| + 1/\tau)^2 + v_F^2 q^2} = \sqrt{W^2 + v_F^2 q^2}, \quad W = |\omega_m| + 1/\tau \quad (29)$$

and

$$S_0(i\omega_m, \mathbf{q}) = \sqrt{|\omega_m|^2 + v_F^2 q^2}. \quad (30)$$

An important feature of the kernel function $K(i\omega_m, \mathbf{q})$, Eq. (28), is that it is exactly zero for $q = 0$ for arbitrary ω_m . Indeed, using $S(i\omega_m, q = 0) = W$, $S_0(i\omega_m, q = 0) = |\omega_m|$, and $\Gamma(i\omega_m, q = 0) = 1/(W\tau - 1)$, we get

$$\begin{aligned} K(i\omega_m, q = 0) &= \frac{1}{W} \left[1 + \frac{1}{W\tau - 1} \right]^2 \left[1 - \frac{1}{W\tau} \right] - \frac{1}{|\omega_m|} \\ &= \frac{1}{W - 1/\tau} - \frac{1}{|\omega_m|} = 0. \end{aligned} \quad (31)$$

We stress that this equality only holds when all the contributions to the self-energy are combined together. This property of the kernel function is characteristic for the gauge-invariant quantities in the presence of interaction. Indeed, the interaction at $q = 0$ implies the shift of the chemical potential and hence can be gauged out². Therefore the contribution of small q to the observables should be suppressed by vanishing of the corresponding kernel function. The same situation is well known for the interaction-induced correction to the conductivity⁹.

It is worth discussing a peculiarity of the problem of magnetooscillations with respect to the gauge invariance. The gauge invariance of the oscillatory part of the observables is guaranteed by the fact that the thermodynamic potential is represented by closed loops. Since the characteristic spatial scale for such a loop is cyclotron radius R_c , the interaction with momenta $q \ll R_c^{-1}$ should not contribute. So, if we would find that our result does not satisfy this requirement, it would mean that the diagrammatic treatment is not sufficient and should be complemented by the infrared cutoff at $q \sim R_c^{-1}$. On a more rigorous level this could be done by using the real-space path integral approach combining the treatment of magnetooscillations in the presence of long-range disorder⁴⁰ with the quantum kinetic-equation approach to the interaction effects^{9,41} and to magnetotransport⁴². In this context, it is instructive to recall the calculation of the dephasing length l_φ , where the infrared cutoff at $q \sim l_\varphi^{-1}$ or $q \sim R^{-1}$ arises for the problems of weak localization⁴³ and Aharonov-Bohm oscillations⁴⁴ in a ring of radius R , respectively.

Since we find, however, that the kernel $K(i\omega_m, q)$ governing the perturbative self-energy (27) does satisfy $K(i\omega_m, q = 0) = 0$, the above cutoff is irrelevant and the perturbative treatment is sufficient in the considered regime of strongly damped oscillations, $T \gg \omega_c$. Indeed, the q -integral in (27) is cut off at $q \sim \omega/v_F \sim T/v_F$ which is much larger than R_c^{-1} under the above condition.

We also emphasize that the kernel $K(i\omega_m, q)$ given by Eq. (28) vanishes in the clean limit

$$K(i\omega_m, q)|_{\tau \rightarrow \infty} = 0. \quad (32)$$

This implies, in particular, that the correction to the effective mass found in Refs. 32,33,45 from the ε - and p -dependence of the self-energy $\Sigma(\varepsilon, p)$ of a clean system, $\delta m/m^* \sim T/E_F$ does not show up in the damping of magnetooscillations, in accordance with the statement made in Ref. 32. In general, $\Sigma(\varepsilon, p)$ is not an observable (and not gauge invariant) quantity, and thus the above correction δm should be at least treated with caution.

III. DAMPING OF MAGNETOOSCILLATIONS: SHORT-RANGE INTERACTION

In this section we evaluate $\delta\Sigma(i\varepsilon_n, \xi_0)$ in the case of a weak point-like interaction given by

$$V(i\omega_m, \mathbf{q}) \equiv U_0. \quad (33)$$

We are interested in the correction to the self-energy to the first order in νU_0 , where $\nu = m^*/2\pi$ is the density of states per spin direction. We calculate the contributions Σ_{ij}^a and Σ_{ij}^b starting from Eqs. (B.15), (B.8), (B.9), (B.20), (B.21), and (B.22) derived in Appendix B. Using the notation $S(i\omega_m, \mathbf{q})$ introduced in Eq. (29), the vertex factor Γ can be presented as

$$\Gamma(i\omega_k, \mathbf{q}) = \frac{1}{S\tau - 1}. \quad (34)$$

Performing the momentum integration for $0 < q < k_F$, we obtain

$$\begin{aligned} \Sigma_{00}^a(i\varepsilon_n, \xi_0) &= -iT U_0 \sum_{\omega_m > \varepsilon_n} \int_0^{k_F} \frac{q dq}{2\pi} \left[\frac{1}{S} - \frac{1}{S_0} \right] \\ &= -\frac{i T U_0}{2\pi v_F^2} \sum_{\omega_m > \varepsilon_n} \left[\sqrt{E_F^2 + W^2} - W - \sqrt{E_F^2 + \omega_m^2} + \omega_m \right] \\ &\simeq -\frac{i T \nu U_0}{2E_F \tau} \sum_{\omega_m > \varepsilon_n} \left[\frac{\omega_m}{\sqrt{E_F^2 + \omega_m^2}} - 1 \right], \end{aligned} \quad (35)$$

$$\begin{aligned} 2\Sigma_{01}^a(i\varepsilon_n, \xi_0) + \Sigma_{11}^a(i\varepsilon_n, \xi_0) &= -\frac{i T U_0}{2\pi v_F^2 \tau} \sum_{\omega_m > \varepsilon_n} \int_{\omega_m + 1/\tau}^{\sqrt{E_F^2 + \omega_m^2}} dS \frac{(2S - 1/\tau)}{(S - 1/\tau)^2} \\ &= -\frac{i T \nu U_0}{2E_F \tau} \sum_{\omega_m > \varepsilon_n} \left(\ln \frac{E_F^2 + \omega_m^2}{\omega_m^2} + \frac{1}{\omega_m \tau} \right). \end{aligned} \quad (36)$$

These expressions are valid independently of the value of the parameter $T\tau$, i.e. they describe both the diffusive and the ballistic regimes, as well as the crossover between them. The logarithmic term in (36) comes from Σ_{01}^a , while the $1/\omega_m$ term originates from Σ_{11}^a .

The contributions of the Hikami-box diagrams are given by

$$\Sigma_{00}^b(i\varepsilon_n) = \frac{i T \nu U_0}{2E_F \tau} \sum_{\omega_m > \varepsilon_n} \left(\omega_m + \frac{1}{\tau} \right) \int_{\omega_m + \frac{1}{\tau}}^{\sqrt{E_F^2 + \omega_m^2}} \frac{dS}{S^2} \simeq -\frac{i T \nu U_0}{2E_F \tau} \sum_{\omega_m > \varepsilon_n} \left[\frac{\omega_m}{\sqrt{E_F^2 + \omega_m^2}} - 1 \right], \quad (37)$$

$$2\Sigma_{01}^b(i\varepsilon_n) + \Sigma_{11}^b(i\varepsilon_n) = \frac{i T U_0}{2\pi v_F^2 \tau^2} \sum_{\omega_m > \varepsilon_n} \left(\omega_m + \frac{1}{\tau} \right) \int_{\omega_m + \frac{1}{\tau}}^{\sqrt{E_F^2 + \omega_m^2}} \frac{dS}{S^2} \frac{(2S - 1/\tau)}{(S - 1/\tau)^2} = \frac{i T \nu U_0}{2E_F \tau} \sum_{\omega_m > \varepsilon_n} \frac{1}{\omega_m \tau}. \quad (38)$$

We see that the Hikami-box contribution exactly cancels the second $(1/\omega_m)$ term in (36). Thus the total correction to the self-energy reads

$$\delta\Sigma(i\varepsilon_n, \xi_0) = -\frac{i T \nu U_0}{E_F \tau} \sum_{\omega_m > \varepsilon_n} \left[\left(\frac{\omega_m}{\sqrt{E_F^2 + \omega_m^2}} - 1 \right) + \ln \frac{\sqrt{E_F^2 + \omega_m^2}}{\omega_m} \right]. \quad (39)$$

The upper limit of the summation over ω_m is effectively given by $m_{\max} \sim E_F/T$. The term $[\omega_m/\sqrt{E_F^2 + \omega_m^2} - 1]$ in Eq. (39), which originates from Σ_{00} , yields after the summation over $\omega_m > \pi T$ a contribution $\propto T$ in addition to a large T -independent contribution, renormalizing τ . As we will see below, a term $\propto T \ln(E_F/T)$ will arise from the contributions of Σ_{01} and Σ_{11} , and hence Σ_{00} yields only a subleading contribution to the damping.

To calculate the sum of logarithms in (39) we write

$$\sum_{\omega_m > \varepsilon_n} \ln \frac{\sqrt{E_F^2 + \omega_m^2}}{\omega_m} = \sum_{\omega_m > \varepsilon_n} \left[\ln \frac{E_F}{\omega_m} + \ln \frac{\sqrt{E_F^2 + \omega_m^2}}{E_F} \right]. \quad (40)$$

For $T \ll E_F$ the second sum can be replaced by the integral, yielding a T -independent contribution. We further use the identity

$$\sum_{m=1}^M \ln \frac{N}{m} = M \ln N - \ln \Gamma(M+1),$$

and the Stirling's formula

$$\ln \Gamma(M+1) = M \ln M - M + \frac{1}{2} \ln(2\pi M) + \dots$$

for $M \gg 1$, where $\Gamma(x)$ is the gamma-function. As a result, we get for arbitrary positive ε_n

$$\delta\Sigma(i\varepsilon_n, \xi_0) \simeq -\frac{iT\nu U_0}{E_F\tau} \left\{ \frac{c_1 E_F}{T} - \frac{\varepsilon_n}{2\pi T} \ln \frac{c_2 E_F}{T} + \ln \left[\frac{1}{\sqrt{2\pi}} \Gamma \left(\frac{\varepsilon_n}{2\pi T} + \frac{1}{2} \right) \right] \right\}, \quad (41)$$

where $c_{1,2}$ are constants of order unity. We see that the contribution to $\delta\Sigma(i\pi T, \xi_0)$ containing vertex corrections to the interaction line has an additional factor $\ln(E_F/T)$ as compared to the T -dependent part of $\delta\Sigma_{00}$. In Eq. (41), we have absorbed the contribution of $\delta\Sigma_{00}$ into the upper cutoff of the log-term which is given by E_F up to a factor of order unity. Furthermore, the same can be done with the last term in Eq. (41) which at $\varepsilon_n = \pi T$ also yields a linear-in- T contribution. Equation (41) thus translates for $T \gg \omega_c$ into the following expression for the damping exponent $B(T) = -2\pi i \delta\Sigma(i\pi T, \xi_0)/\omega_c$:

$$B(T) = -c_1 \nu U_0 \frac{\pi}{\omega_c \tau} + \frac{\pi T}{\omega_c} \frac{\nu U_0}{E_F \tau} \ln \frac{c_2 E_F}{T}. \quad (42)$$

The first term in Eq. (42) describes the T -independent FL-renormalization of τ due to vertex corrections and should be included in the effective relaxation time τ^* , as was mentioned after Eq. (15) in Sec. II. The second term represents the T -dependent contribution to the damping factor that we are interested in and is analyzed in the next section.

IV. INTERPRETATION: EFFECTIVE MASS VS QUANTUM SCATTERING TIME

The above result (42) can be interpreted in terms of corrections to the effective mass (or ω_c) and the elastic scattering rate entering the standard formula (17). These corrections come from the interplay of disorder and interaction. Writing

$$\begin{aligned} A_1(T) &= \frac{4\pi^2 T}{\omega_c} \exp \left[-\frac{2\pi^2 T}{(\omega_c + \delta\omega_c)} - \frac{\pi}{(\omega_c + \delta\omega_c)} \frac{1}{(\tau + \delta\tau)} \right] \\ &\simeq A_1^{(0)}(T) \exp \left[\frac{2\pi^2 T}{\omega_c} \frac{\delta\omega_c}{\omega_c} \right] \exp \left[\frac{\pi}{\omega_c \tau} \left(\frac{\delta\omega_c}{\omega_c} + \frac{\delta\tau}{\tau} \right) \right] \\ &\simeq A_1^{(0)}(T) \exp \left[-\frac{2\pi^2 T}{\omega_c} \frac{\delta m}{m} \right] \exp \left[\frac{\pi}{\omega_c \tau} \left(-\frac{\delta m}{m} + \frac{\delta\tau}{\tau} \right) \right], \end{aligned} \quad (43)$$

we conclude that

$$B(T) = -\frac{2\pi^2 T}{\omega_c} \frac{\delta m}{m} - \frac{\pi}{\omega_c \tau} \left(\frac{\delta m}{m} - \frac{\delta\tau}{\tau} \right). \quad (44)$$

It is worth noting that the FL-renormalization does not affect the product $\omega_c m = eB$.

Comparing (42) and (44) [we recall that the first term in Eq. (42) is absorbed in τ], we see that the $T \ln T$ dependence of the damping factor could in principle originate either from the $\ln T$ correction to the effective mass, or from the $T \ln T$ -type correction to τ . This led the authors of Ref. 26 to the conclusion that the nonlinear T -dependence of the damping factor may be equivalently interpreted either as a T -dependent renormalization of the effective mass or as a T -dependent Dingle temperature. It is clear, however, that these two possibilities correspond to different physical processes.

A. Self-energy at real energies: analytical continuation

To identify the physical origin of the leading contribution to the damping it is instructive to obtain $B(T)$ using the expression for the self-energy analytically continued to real values of energies $\varepsilon_n \rightarrow -i\varepsilon$. Performing the analytical continuation to real energies ε and real frequencies ω in Eq. (27), we get

$$\delta\Sigma(\varepsilon, \xi_0) = \frac{\nu U_0}{E_F \tau} \int_{-\infty}^{\infty} \frac{d\omega}{4\pi} \tanh\left(\frac{\varepsilon - \omega}{2T}\right) \left\{ \ln\left(\frac{\sqrt{E_F^2 - \omega^2}}{-i\omega}\right) + \left(\frac{-i\omega}{\sqrt{E_F^2 - \omega^2}} - 1\right) \right\}, \quad (45)$$

so that the real part of the self-energy is given by

$$\begin{aligned} \text{Re } \delta\Sigma(\varepsilon, \xi_0) &= \frac{\nu U_0}{4\pi E_F \tau} \left\{ \int_0^{E_F} d\omega \left[\tanh\left(\frac{\varepsilon - \omega}{2T}\right) + \tanh\left(\frac{\varepsilon + \omega}{2T}\right) \right] \left[\ln\left(\frac{\sqrt{E_F^2 - \omega^2}}{\omega}\right) - 1 \right] \right. \\ &\quad \left. + \int_{E_F}^{\infty} d\omega \left[\tanh\left(\frac{\varepsilon - \omega}{2T}\right) + \tanh\left(\frac{\varepsilon + \omega}{2T}\right) \right] \left[\ln\left(\frac{\sqrt{\omega^2 - E_F^2}}{\omega}\right) + \frac{\omega}{\sqrt{\omega^2 - E_F^2}} - 1 \right] \right\} \\ &\simeq \varepsilon \frac{\nu U_0}{2\pi E_F \tau} \ln \frac{E_F}{\max[|\varepsilon|, T]}. \end{aligned} \quad (46)$$

The leading contribution here comes from the term $\ln(E_F/\omega)$ in the first integral over $\omega < E_F$ while other terms only rescale the ultraviolet cut-off E_F of the logarithm by a constant of order unity, which is beyond the accuracy of our quasichlassical approximation.

The imaginary part of $\delta\Sigma$ reads

$$\begin{aligned} \text{Im } \delta\Sigma(\varepsilon, \xi_0) &= \frac{\nu U_0}{4\pi E_F \tau} \int_0^{E_F} d\omega \left[\tanh\left(\frac{\varepsilon - \omega}{2T}\right) - \tanh\left(\frac{\varepsilon + \omega}{2T}\right) \right] \left[\frac{\pi}{2} - \frac{\omega}{\sqrt{E_F^2 - \omega^2}} \right] \\ &\simeq -\text{const} \frac{\nu U_0}{\tau} + \frac{\nu U_0}{2E_F \tau} T \ln \left[2 \cosh\left(\frac{\varepsilon}{2T}\right) \right], \end{aligned} \quad (47)$$

where the T -dependent term has the following asymptotics:

$$T \ln[2 \cosh(\varepsilon/2T)] = \begin{cases} \varepsilon/2, & \varepsilon \gg T, \\ T \ln 2, & \varepsilon \ll T. \end{cases} \quad (48)$$

The contribution of the term $\omega/\sqrt{E_F^2 - \omega^2}$ to the integral in (47) is T -independent up to small corrections of order of $\nu U_0(T/E_F)^2/\tau$ which are beyond the accuracy of the calculation.

Having calculated $\text{Re } \Sigma$ and $\text{Im } \Sigma$ for real energies ε , we can determine δm and $\delta\tau$. Indeed, the magnitude of the first harmonics of the magnetooscillations of the thermodynamic density of states is expressed through the real- ε self-energy $\delta\Sigma(\varepsilon)$ as follows:

$$A_1(T) = \frac{4\pi^2 T}{\omega_c} \int d\varepsilon \left[-\frac{\partial n_F(\varepsilon)}{\partial \varepsilon} \right] A_1(\varepsilon, T), \quad (49)$$

$$A_1(\varepsilon, T) = \exp \left\{ \frac{2\pi i}{\omega_c} [\varepsilon - \text{Re } \delta\Sigma(\varepsilon, \xi_0)] \right\} \exp \left\{ -\frac{\pi}{\omega_c \tau} + \frac{2\pi}{\omega_c} \text{Im } \delta\Sigma(\varepsilon, \xi_0) \right\}, \quad (50)$$

where $n_F(\varepsilon) = [1 + \exp(\varepsilon/T)]^{-1}$ is the Fermi distribution function.

In analogy with Eq. (43) we represent the energy-dependent amplitude $A_1(\varepsilon, T)$ in terms of energy- and temperature-dependent corrections to the quantum scattering time and mass, $\delta\tau(\varepsilon, T)$ and $m(\varepsilon, T)$:

$$A_1(\varepsilon, T) = \exp \left\{ \frac{2\pi i \varepsilon}{\omega_c} \left[1 + \frac{\delta m(\varepsilon, T)}{m} \right] \right\} \exp \left\{ -\frac{\pi}{\omega_c \tau} \left[1 + \frac{\delta m(\varepsilon, T)}{m} - \frac{\delta\tau(\varepsilon, T)}{\tau} \right] \right\}. \quad (51)$$

Comparing (51) with (50), we express $\delta\tau(\varepsilon, T)$ and $m(\varepsilon, T)$ through $\text{Re } \Sigma(\varepsilon)$ and $\text{Im } \Sigma(\varepsilon)$ as follows:

$$\frac{\delta m(\varepsilon, T)}{m} = -\frac{\text{Re } \delta\Sigma(\varepsilon, T)}{\varepsilon}, \quad (52)$$

$$\frac{\delta\tau(\varepsilon, T)}{\tau} = 2\tau \text{Im } \delta\Sigma(\varepsilon, T) + \frac{\delta m(\varepsilon, T)}{m}. \quad (53)$$

Using (46) and (47) in combination with (52) and (53), we obtain

$$\frac{\delta m(\varepsilon, T)}{m} = -\frac{\nu U_0}{2\pi E_F \tau} \ln \frac{E_F}{\max[|\varepsilon|, T]}, \quad (54)$$

$$\frac{\delta \tau(\varepsilon, T)}{\tau} = \nu U_0 \frac{T}{E_F} \ln \left[2 \cosh \left(\frac{\varepsilon}{2T} \right) \right] - \frac{\nu U_0}{2\pi E_F \tau} \ln \frac{E_F}{\max[|\varepsilon|, T]}. \quad (55)$$

The integration in (49) sets in effect $\varepsilon \sim T$ in the above expressions. The T -dependent corrections to the effective mass and the quantum scattering time extracted experimentally with the help of Eq. (43) are thus given by

$$\frac{\delta m(T)}{m} = -\frac{\nu U_0}{2\pi E_F \tau} \ln \frac{E_F}{T}, \quad (56)$$

$$\frac{\delta \tau(T)}{\tau} = \nu U_0 \frac{T}{E_F} - \frac{\nu U_0}{2\pi E_F \tau} \ln \frac{E_F}{T}. \quad (57)$$

In (57) we assume that $\pi \nu U_0 T / E_F \ll \omega_c \tau$, expand $\exp[\pi \delta \tau(\varepsilon, T) / \omega_c \tau^2]$, and then average the term $\ln[2 \cosh(\varepsilon/2T)]$ [which is a real-energy counterpart of the last term in Eq. (41)] with $-\partial n_F(\varepsilon) / \partial \varepsilon$.

It is clear from these results that the leading term in $B(T)$ [proportional to $T \ln(E_F/T)$, Eq. (42)] originates from the real part of the self-energy, Eq. (46), i.e. from renormalization of the effective mass, which affects incommensurability of the oscillations at different values of energy ε . The contribution to $B(T)$ of the imaginary part of the self-energy [corresponding to the last term in Eq. (41)], which is governed in the ballistic regime by the renormalization of the scattering time, is smaller by a factor $\ln(E_F/T)$. In the expression for the damping, Eq. (42), this contribution is absorbed in the numerical constant c_2 in the upper cutoff of the logarithm.

The obtained result for the interaction-induced correction to the scattering time τ , Eq. (57), agrees, up to a factor $\frac{1}{2}$, with the correction to the transport time following from the calculation of conductivity correction in the ballistic regime in Ref. 9. This is exactly what one would expect on physical grounds. Indeed, it is known that the conductivity correction⁹ can be understood as governed by an additional, predominantly back-scattering, contribution to the scattering cross-section related to the dressing of an impurity by Friedel oscillations. Since this contribution is concentrated near the scattering angle $\phi = \pi$, the correction to the momentum relaxation rate is larger by the factor $1 - \cos \phi \simeq 2$ than the correction to the total scattering rate. In Sec. IV B we will corroborate the results of this subsection by an explicit calculation of the contribution to the impurity scattering rate due to Friedel oscillations.

Up to now we calculated the exchange contribution to the self-energy. For the point-like interaction, the Hartree term has opposite sign and is twice larger in magnitude than the exchange term due to the spin summation. This simply reverses the sign of the corrections to the damping factor.

B. Calculation of $\delta \tau$ from the scattering off Friedel oscillations.

In this subsection we calculate the correction to the total elastic scattering time $\delta \tau$ in a different, physically more transparent way, considering the scattering off impurities dressed by Friedel oscillations^{9,38}. We will demonstrate how the result (57) is reproduced in this way. In particular, this will confirm once more that there is no $T \ln T$ term in $\delta \tau$ and therefore the leading $T \ln T$ contribution to the damping factor comes from δm .

We start with the expression relating the total elastic scattering rate and the scattering cross-section $S(\phi)$ of a single impurity,

$$\frac{1}{\tau_q(\varepsilon, T)} = n_{\text{imp}} v_F \int \frac{d\phi}{2\pi} S(\phi), \quad (58)$$

where n_{imp} is the concentration of impurities. The expression for the transport scattering time τ_{tr} determining the conductivity differs from (58) by a factor $1 - \cos \phi$ in the integrand:

$$\frac{1}{\tau_{\text{tr}}(\varepsilon, T)} = n_{\text{imp}} v_F \int \frac{d\phi}{2\pi} S(\phi) (1 - \cos \phi). \quad (59)$$

We note that the two times, τ_q and τ_{tr} , though equal for the point-like impurities in the non-interacting case [for which $S(\phi) = (n_{\text{imp}} v_F \tau)^{-1} = \text{const}(\phi)$], differ from each other when the scattering off Friedel oscillations is taken into account.

The impurity scattering cross-section for dressed impurities reads

$$S(\phi) = \frac{2\pi\nu}{v_F} \left| V \left(2k \sin \frac{\phi}{2} \right) \right|^2 \simeq S_0 + \frac{2\pi\nu V_0}{v_F} 2 \text{Re} \delta V \left(2k \sin \frac{\phi}{2} \right). \quad (60)$$




FIG. 4: Schematic illustration of the scattering off Friedel oscillations. The black dot in the middle represents a short-range impurity which creates the oscillatory correction to the electron density around it. The circle represents the equipotential line of the effective impurity potential. The correction to the impurity cross-section at the angle ϕ arises due to the interference of two electronic waves^{9,38}, one of which (dashed line) scattered by the impurity and another (solid line) by the Friedel oscillations at a point parametrized by the distance r from the impurity and the angle ψ .

Here $S_0 = 2\pi\nu V_0^2/v_F$ is the bare impurity scattering cross-section and $V_0 = \int d^2r V_0(r)$, where $V_0(r)$ is the bare point-like impurity potential. The cross-section $S(\phi)$ depends on energy ϵ of an electron through $k = k_F + \epsilon/v_F$ in the Fourier transform $V(q) = V_0 + \delta V(q)$ of the effective impurity potential, renormalized by the Friedel oscillations of the electron density. For $r \gg k_F^{-1}$ the oscillatory correction to the electron density reads

$$\delta\rho(r) = -\nu V_0 \frac{(2\pi r T/v_F)}{\sinh(2\pi r T/v_F)} \frac{\sin 2k_F r}{\pi r^2}. \quad (61)$$

The correction to the impurity scattering potential $\delta V(r)$ due to scattering off the Friedel oscillations is proportional, for the short-range interaction, to the electron density at point $\mathbf{r}$. Similarly to the consideration of Sec. III and Sec. IV A, we will concentrate on the exchange part of this correction,

$$\delta V(r) = -\frac{1}{2} U_0 \delta\rho(r). \quad (62)$$

To calculate the correction to the impurity cross-section, we need the Fourier transform of $\delta V(r)$,

$$\delta V\left(2k \sin \frac{\phi}{2}\right) = \frac{U_0}{2} \int d^2r \delta\rho(r) \exp\left(2i k r \sin \frac{\phi}{2} \cos \psi\right), \quad (63)$$

where ψ is the polar angle of $\mathbf{r}$, see Fig. 4. Substituting (61) and (63) into (60), we find the interaction-induced correction to the scattering cross-section,

$$\delta S(\phi) = \nu U_0 S_0 \int \frac{d\psi}{2\pi} \int dr \frac{(2\pi T/v_F)}{\sinh(2\pi r T/v_F)} \left\{ \sin\left(2r \left[k_F + k \cos \psi \sin \frac{\phi}{2}\right]\right) + \sin\left(2r \left[k_F - k \cos \psi \sin \frac{\phi}{2}\right]\right) \right\}. \quad (64)$$

Performing the integration over r we get

$$\delta S(\phi) = S_0 \frac{\nu U_0}{4} \int d\psi [\tanh \beta_+ + \tanh \beta_-], \quad (65)$$

where

$$\beta_{\pm} = \frac{v_F}{2T} \left(k_F \pm k \cos \psi \sin \frac{\phi}{2}\right) = \frac{E_F}{T} \left[1 \pm \left(1 + \frac{\epsilon}{2E_F}\right) \cos \psi \sin \frac{\phi}{2}\right]. \quad (66)$$

For $\beta_{\pm} \gg 1$, i.e. for most values of the scattering angle ϕ except for those corresponding to the backscattering ($\phi \approx \pi$), we see that the scattering cross-section does not depend on ϕ up to exponentially small corrections of order $\mathcal{O}(\exp[-E_F/T])$:

$$S(\phi) = S_0 [1 + \pi\nu U_0], \quad |\phi - \pi| \gg (\max[T, \epsilon]/E_F)^{1/2}. \quad (67)$$




FIG. 5: Differential impurity cross-section $S(\phi)$ (in units of the bare cross-section S_0) renormalized by the scattering off Friedel oscillations (with only the exchange contribution taken into account) calculated for $\nu U_0 = 0.2$ and (a) fixed $T = 0.01E_F$ and $\epsilon/E_F = 0, 0.01, 0.02, 0.05, 0.1, 0.2$, from top to bottom; (b) fixed $\epsilon = 0$ and $T/E_F = 0.005, 0.01, 0.02, 0.05, 0.1, 0.2$, from top to bottom.

On the other hand, in the vicinity of $\phi = \pi$ the cross-section has a “hump”, see Fig. 5, of the width and the height scaling as $\delta\phi$, $\delta S(\phi)/S_0 \sim (\max[T, \epsilon]/E_F)^{1/2}$. The explicit expression for $S(\phi)$ at $T = 0$ can be found in Ref. 9. In Fig. 5 we plotted $S(\phi)$ for fixed T and several values of ϵ (Fig. 5a) and for fixed $\epsilon = 0$ and several values of T (Fig. 5b).

To calculate the (ϵ, T) -dependent correction to $1/\tau_q$ (which is determined by the “hump”), we expand $\sin(\phi/2)$ around $\phi = \pi$. Furthermore, we expand $\cos\psi$ around $\psi = \pi$ and $\psi = 0$ in the expressions for β_+ and β_- , respectively. This corresponds to the two possible interfering paths propagating along the horizontal line in Fig. 4 with the scattering off the Friedel oscillation occurring either to the left or to the right from the impurity. Denoting $\delta\psi = x$, $\delta\phi/2 = y$, $z^2 = x^2 + y^2$, and $\omega = z^2 E_F$, we get from Eq. (65)

$$\begin{aligned} \int \frac{d\phi}{2\pi} \delta S(\phi) &\simeq S_0 \frac{\nu U_0}{2\pi} \int dx \int dy \tanh \left\{ \frac{E_F}{T} \left[1 - \left(1 + \frac{\epsilon}{2E_F} \right) \left(1 - \frac{x^2 + y^2}{2} \right) \right] \right\} \\ &\simeq S_0 \nu U_0 \int z dz \tanh \left\{ \frac{E_F}{T} \left[\frac{z^2}{2} - \frac{\epsilon}{2E_F} \right] \right\} = S_0 \frac{\nu U_0}{2E_F} \int_0^{\sim E_F} d\omega \tanh \left(\frac{\omega - \epsilon}{2T} \right) \end{aligned} \quad (68)$$

Using (58) and (68), we find the total quantum scattering rate,

$$\frac{1}{\tau_q(\epsilon, T)} = \frac{1}{\tau} + \delta \left(\frac{1}{\tau_q} \right), \quad (69)$$

$$\delta \left(\frac{1}{\tau_q} \right) = -\frac{\delta \tau_q(\epsilon, T)}{\tau^2} = \frac{\nu U_0}{\tau} \left[\text{const} - \frac{T}{E_F} \ln \left(2 \cosh \frac{\epsilon}{2T} \right) - \frac{\epsilon}{2E_F} \right]. \quad (70)$$

This result for the interaction-induced correction to τ_q agrees (to the leading order in $1/T\tau$ corresponding to the ballistic regime) with that obtained from the imaginary part of the self-energy, Eqs. (55) and (57). More accurately, Eq. (70) differs from Eq. (55) by the last (T -independent) term $\nu U_0 \epsilon / 2E_F \tau$, which drops out after the thermal averaging with $-\partial n_F(\epsilon)/\partial \epsilon$ and thus does not contribute to $\delta \tau_q(T)$, Eq. (57). This term is in a sense anomalous, since it arises from the ultraviolet limit of the ω -integration in Eq. (68). In fact, one could question the validity of this contribution, since we used the asymptotic, large- r form of the Friedel oscillations in Eq. (61). One can check, however, that the same result [up to an irrelevant additive constant independent of T and ϵ : in Eq. (67) $\pi \nu U_0$ is replaced by $2\nu U_0$] is obtained from a calculation using the exact form of the Friedel oscillations⁴⁶. The appearance of this linear-in- ϵ term is related to the violation of the particle-hole symmetry in the parabolic spectrum; this term did not appear in the diagrammatic calculation of Sec. IV where the spectrum was linearized. What however enters the experimental damping of magnetooscillations is $1/\tau_q$ integrated over the energy with an even function $-\partial n_F(\epsilon)/\partial \epsilon$. Therefore, we are in fact interested in the collision rate symmetrized with respect to $\epsilon \rightarrow -\epsilon$. Performing this symmetrization in Eq. (64), we get a result determined solely by the infrared scale, $r \sim v_F/T$, yielding Eq. (70) without the last, linear-in- ϵ term.

Finally, using (59), we see that the correction to the transport rate is larger than $\delta(1/\tau_q)$ by a factor of 2. The corresponding correction to the conductivity reproduces the result of Ref. 9 in the ballistic limit.




FIG. 6: Diagrammatic equation for the effective interaction line (bold wavy line) in the random-phase approximation, Eqs. (71) and (73). Dotted wavy line represents the bare interaction, $V_0(q) + F_0^\rho/2\nu$ in the singlet channel or $F_0^\sigma/2\nu$ in the triplet channel. The bubble Π is the polarization operator, Eq. (72).

V. DAMPING OF MAGNETOOSCILLATIONS: COULOMB INTERACTION

We turn now to the Coulomb interaction. In the case of Coulomb interaction, one should take into account the dynamical screening of the interaction within the random phase approximation (RPA), see Fig. 6. In what follows we use for simplicity the so-called F_0 -approximation⁹, which retains only the zeroth harmonics of the Fermi-liquid constants F_0^ρ and F_0^σ in the charge and spin channels, respectively. Then the effective interaction propagator in the charge channel (combining the exchange term and the singlet contribution of the Hartree-type interaction) reads^{9,10}

$$V^\rho(i\omega_m, q) = [(V_0(q) + F_0^\rho/2\nu)^{-1} + \Pi(i\omega_m, q)]^{-1}, \quad (71)$$

where $\Pi(i\omega_m, q)$ is the polarization operator

$$\Pi(i\omega_m, q) = 2\nu [1 - |\omega_m| \tau \Gamma(i\omega_m, q)], \quad (72)$$

$\Gamma(i\omega_m, q)$ is the impurity ladder, and $\nu = m/2\pi$ is the density of states per spin direction. The triplet contribution to the effective interaction arises from the ladder of Hartree-type interaction blocks and reads

$$V^\sigma(i\omega_m, q) = [2\nu/F_0^\sigma + \Pi(i\omega_m, q)]^{-1}. \quad (73)$$

A. Singlet channel

The main difference as compared to the case of the short-range weak interaction considered in Sec. III is the nontrivial form of the dynamically screened Coulomb interaction (one should take into account the renormalization of the interaction by polarization operator). Using

$$V_0(q) = \frac{2\pi e^2}{q}$$

and neglecting F_ρ^0 , we get

$$\begin{aligned} V^\rho(i\omega_m, q) &= \frac{V_0(q)}{1 + V_0(q)\Pi(i\omega_m, q)} = \frac{2\pi e^2/q}{1 + (2\pi e^2/q)2\nu[1 - |\omega_m|\tau\Gamma(i\omega_m, q)]} \\ &= \frac{1}{2\nu} \frac{\kappa(S - 1/\tau)}{(q + \kappa)(S - 1/\tau) - \kappa(W - 1/\tau)}. \end{aligned} \quad (74)$$

Here we use the standard notation $\kappa = 4\pi\nu e^2$ for the inverse Thomas–Fermi screening radius and use the short-hand notation S for $S(i\omega_m, \mathbf{q}) \equiv \sqrt{W^2 + v_F^2 q^2}$ with $W = |\omega_m| + 1/\tau$.

For $q \ll \kappa$, neglecting q in the sum $q + \kappa$ in the denominator of (74), one finds that the exchange interaction

$$\tilde{V}^\rho(i\omega_m, q) = \frac{1}{2\nu} \frac{S - 1/\tau}{S - W} \quad (75)$$

has a singularity $\propto 1/q^2$ in the limit $q \rightarrow 0$:

$$\tilde{V}^\rho(i\omega_m, q) = \frac{1}{2\nu} \frac{2|\omega_m|(|\omega_m| + 1/\tau)}{q^2 v_F^2}, \quad q \rightarrow 0, \quad (76)$$

so that each separate term $\Sigma_{ij}^a, \Sigma_{ij}^b$ in the self-energy would diverge. This divergence is analogous to the one encountered in course of calculation of the tunneling density of states¹. In that case, one has to keep the q term

in denominator of (74), which cuts off the logarithmic divergence. For the present problem, this is, however, not needed. Indeed, as was emphasized in the end of Sec. II, the kernel $K(i\omega_m, q)$ combining together contributions of all relevant self-energy diagrams is proportional to q^2 in the limit $q \rightarrow 0$ and hence cancels the singularity in $\tilde{V}(i\omega_m, q)$. In view of this, it is convenient to represent the kernel function (28) in a form which shows explicitly that $K(i\omega_m, q = 0) = 0$ [we recall that $S_0(q = 0) = |\omega_m|$ and $S(q = 0) = W$]:

$$K(i\omega_m, q) = \left(-\frac{1}{S_0} + \frac{1}{|\omega_m|} \right) + \frac{S - W}{S - 1/\tau} \left[\frac{1}{S\tau(S - 1/\tau)} - \frac{1}{|\omega_m|} \right]. \quad (77)$$

Then the product $\tilde{V}^\rho(i\omega_m, q)K(i\omega_m, q)$ takes the form

$$2\nu\tilde{V}^\rho(i\omega_m, q)K(i\omega_m, q) = \frac{S_0 - |\omega_m|}{S_0(S - W)} - \frac{1}{S_0} + \frac{1}{S\tau(S - 1/\tau)}. \quad (78)$$

Performing the integration over the momentum q in (27), we get the correction to self-energy in the singlet (charge) channel

$$\delta\Sigma^\rho(i\varepsilon_n, \xi_0) = -\frac{i T}{2E_F\tau} \sum_{\omega_m > \varepsilon_n}^{\Delta} \left\{ (1 + 2\omega_m\tau) \ln \frac{1 + 2\omega_m\tau}{2\omega_m\tau} - 1 + \ln \frac{\Delta^2 + \omega_m^2}{\omega_m^2} \right\}, \quad (79)$$

where we introduced

$$\Delta \equiv \kappa v_F. \quad (80)$$

Comparing (79) and (39), we see that the last term in (79) corresponds to a static short-range interaction with $\nu U_0 = 1$.

Setting $\varepsilon_n = \pi T$ and separating the contributions to the sum (79) governed by the high-energy ($\varepsilon \sim \Delta$) and low-energy ($\varepsilon \sim T$) regions, we can present the result in the following form:

$$\delta\Sigma^\rho(i\pi T, \xi_0) = -\frac{i T}{2E_F\tau} \left[\frac{\text{const}\Delta}{T} - \left(1 - \frac{1}{8\pi T\tau} \right) \ln \frac{\Delta}{T} - f(4\pi T\tau) \right], \quad (81)$$

where $f(x)$ is a parameterless function,

$$f(x) = \sum_{m=1}^{\infty} \left[1 - (1 + mx) \ln \frac{1 + mx}{mx} + \frac{1}{2mx} \right] = \begin{cases} \frac{c_f}{x} + \left[\frac{1}{2x} + \frac{1}{2} \right] \ln \frac{1}{x}, & x \ll 1, \\ \frac{\pi^2}{36 x^2}, & x \gg 1, \end{cases} \quad (82)$$

with $c_f = -3/4 - \psi(1)/2 = -0.461392..$ (here $\psi(x)$ is the digamma function). Thus the dynamical screening of Coulomb interaction leads to different asymptotics of the self-energy in the diffusive and ballistic regimes, in contrast to the case of weak short-range interaction⁴⁷.

The T -dependence of the leading correction to the magnetooscillations damping factor due to the interaction in the singlet channel has therefore the form

$$B^\rho(T) = \frac{\pi}{\omega_c\tau} \frac{T}{E_F} \left[\left(1 - \frac{1}{8\pi T\tau} \right) \ln \frac{\Delta}{T} + f(4\pi T\tau) \right] \quad (83)$$

$$= \frac{\pi}{\omega_c\tau} \frac{T}{E_F} \times \begin{cases} \frac{3}{2} \ln \frac{\Delta}{T} - \frac{1}{2} \left(1 + \frac{1}{4\pi T\tau} \right) \ln(4\pi\Delta\tau) - \frac{c_f}{4\pi T\tau}, & 4\pi T\tau \ll 1, \\ \left(1 - \frac{1}{8\pi T\tau} \right) \ln \frac{\Delta}{T}, & 4\pi T\tau \gg 1. \end{cases} \quad (84)$$

This result is illustrated in Fig. 7. Note that the ballistic asymptotics describes the exact result with a remarkable accuracy down to very low temperature, $T\tau \sim 0.01 - 0.05$, see Fig. 7a. Retaining in Eq. (84) only the leading terms and suppressing the T -independent contributions which can be absorbed in the FL-renormalized τ , we get

$$B^\rho(T) = \frac{\pi}{\omega_c\tau} \frac{T}{E_F} \times \begin{cases} \frac{3}{2} \ln \frac{\Delta}{T} - \frac{1}{2} \ln(4\pi\Delta\tau), & 4\pi T\tau \ll 1, \\ \ln \frac{\Delta}{T}, & 4\pi T\tau \gg 1. \end{cases} \quad (85)$$




FIG. 7: Temperature dependence of the singlet channel correction to the damping factor $B^\rho(T)$, Eq. (83), for $4\pi\Delta\tau = 100$ (solid line) with the low- T (dot-dashed) and high- T (dashed) asymptotics, Eq. (84). (a) Wide temperature range: on this scale $B^\rho(T)$ is essentially indistinguishable from its high- T asymptotics; (b) low- T part: the crossover between the two asymptotics occurs at $T\tau \sim 0.05$.

B. Triplet channel

Calculation of the corresponding triplet contribution $B^\sigma(T)$ is presented in Appendix C and leads to qualitatively similar asymptotics. The leading term in the total correction to the damping factor in the ballistic regime, realized in experiments on low-disorder samples at realistic temperatures, takes the simple form

$$B(T) = B^\rho(T) + B^\sigma(T) \simeq \left(1 + \frac{3F_0^\sigma}{1 + F_0^\sigma}\right) \frac{\pi}{\omega_c\tau} \frac{T}{E_F} \ln \frac{\Delta}{T}. \quad (86)$$

As discussed in Sec. IV, this result arises due to the correction to the effective mass (the consideration of Sec. IV fully applies to the case of the Coulomb interaction as well).

It is worth noting that due to the factor $(1 + F_0^\sigma)$ in the denominator of $B^\sigma(T)$, the damping of magnetooscillations tends to diverge upon approaching the Stoner ferromagnetic instability. Since the damping is determined by the effective mass $m^*(T)$ (see Sec. IV), we conclude that $m^*(T)$ diverges as $F_0^\sigma \rightarrow -1$ due to the interplay of disorder and interaction. This should be contrasted with the clean case, where the effective mass is solely determined by Fermi-liquid constant in the singlet channel, F_1^ρ , and hence is insensitive to the ferromagnetic instability.

C. Discussion and comparison to earlier work

Let us discuss the obtained results for the damping factor $B(T)$. In both the diffusive and ballistic regimes we find the temperature dependence of the form $B(T) \propto T \ln T$. In the diffusive (low- T) regime the relative correction to the damping factor, $\omega_c\tau B \sim g^{-1}T\tau \ln T$ is less singular than the known corrections¹ to the conductivity, $\delta\sigma/\sigma \sim g^{-1} \ln T$, and the tunneling density of states, $\delta\rho/\rho \sim g^{-1} \ln^2 T$ (here $g \sim E_F\tau$ is the dimensionless conductivity). On the technical level, the qualitative difference with the tunneling DOS can be traced back to the contribution of the Hikami box diagrams, Σ_{ij}^b .

As has been already mentioned in Sec. I, the problem of the effect of the interaction on magnetooscillations was recently addressed in Ref. 26. The result of this work for the damping factor is qualitatively similar to ours, $B \propto T \ln T$. However, the crossover function and, in particular, the prefactors in both ballistic and diffusive limits differ from ours (1 instead of 3/2 in the first line of Eq. (85), and 3/2 instead of 1 in the second line). This difference is not surprising, since the authors of Ref. 26 took into account only one diagram Σ_{01}^a out of six in Fig. 2. Thus, even the qualitative agreement may be considered as an accidental coincidence. In fact, there is a conceptual difference between our result and that of Ref. 26. To illustrate this, consider a toy interaction of the type $V(\omega, \mathbf{q}) = F(\omega)\delta(\mathbf{q})$. Our result then would be zero, since the kernel function $K(i\omega_m, \mathbf{q})$, Eq. (28), satisfies the gauge-invariance constraint [see Eq. (31)]

and discussion below it],

$$K(i\omega_m, \mathbf{q})|_{\mathbf{q}=0} = 0. \quad (87)$$

In contrast, the formula of Ref. 26 would give a finite result, since their kernel

$$K_{01}^{(a)} = \frac{2}{S} \frac{1}{S\tau - 1} \quad (88)$$

does not satisfy the requirement (87),

$$K_{01}^{(a)}(i\omega_m, q=0) = \frac{2}{|\omega_m|(|\omega_m|\tau + 1)}.$$

For the Coulomb interaction this results in a logarithmic divergency at small q that is cut off by the plasmon pole at $q \sim q_{\min}$ with

$$q_{\min} = \frac{|\omega_m|(|\omega_m| + 1/\tau)}{\kappa v_F^2},$$

similarly to the calculation of the tunneling density of states¹. As we explained in the end of Sec. II, the contribution of small momenta, $q < R_c^{-1}$, should be suppressed for the present problem. Therefore, the above small- q divergence should be cut off by the magnetic field, which would partly transform $\ln T$ of Ref. 26 into $\ln B$. However, this problem is in fact spurious: the result of our work does not suffer from any infrared divergencies, since our kernel $K(i\omega_m, \mathbf{q})$ does satisfy Eq. (87).

Finally, let us briefly comment on the ultra-diffusive regime, $T \ll \omega_c \ll 1/\tau$. In this regime the summation over Matsubara energies ε_n is not restricted to $n = 0$ and $N_c \sim \omega_c/T \gg 1$ Matsubara harmonics are important. Therefore, the damping due to the inelastic scattering [suppressed only for $n = 0$, see Ref. 26 and Eq. (B.13)] becomes finite. The corresponding contribution to the damping can be roughly estimated using Eq. (B.13) taken at relevant $n \sim N_c$:

$$\delta B^{\text{inel}}(T) \sim -n^2 \frac{T^2}{\omega_c E_F} \ln(E_F \tau) \Big|_{n \sim N_c} \sim -\frac{\omega_c}{E_F} \ln(E_F \tau), \quad (89)$$

yielding $|\delta B^{\text{inel}}(T)| \ll 1$, since in the ultra-diffusive regime $\omega_c \ll 1/\tau$. Thus the inelastic contribution to the damping factor is always small. Note that the contribution to the damping due to the renormalization of the effective mass in the ultra-diffusive regime is also small: at $n = N_c$ we have $B(T) = -2\pi\varepsilon_n/\omega_c \delta m/m \sim \delta m/m \ll 1$.

VI. CONCLUSIONS

In conclusion, we have studied the T -dependent interaction corrections to the damping of magnetooscillations in a two-dimensional electron gas. The damping factor has been calculated for Coulomb and short-range interaction in the whole range of temperatures, from the ballistic to the diffusive regime. While the relevant diagrams are similar to those for the local density of states, the results are essentially different, see Eqs. (42), (83).

We have identified leading contributions to the damping induced by interplay of interaction and disorder, which can be associated with corrections to the effective mass and the quantum scattering time. It has been shown that in the ballistic regime, which is typically realized in low-disorder samples at realistic temperatures, the dominant effect is that of the renormalization of the effective electron mass due to the interplay of the interaction and impurity scattering. Specifically, the correction to the effective mass is of the form $\delta m/m \sim 1/(E_F \tau) \ln(E_F/T)$, Eq.(56). The correction to the impurity scattering time is of the form $\delta \tau_q/\tau \sim T/E_F$, Eq. (57), and yields a subleading contribution to the damping. We have confirmed the result for the correction to the quantum scattering time by performing a calculation based on the picture of scattering by impurities dressed by Friedel oscillations. The results of the paper are relevant to the analysis of experiments on magnetooscillations (in particular, for extracting the value of the effective mass) and are expected to be useful for understanding the physics of a high-mobility 2DEG near the apparent metal-insulator transition.

VII. ACKNOWLEDGMENTS

We thank I. Aleiner, D. Aristov, A. Chubukov, S. Kravchenko, and V. Pudalov for useful discussions. We are particularly grateful to D. Maslov and M. Reizer for discussions of Ref. 26 and comments on the draft of our paper. This work was supported by SPP ‘‘Quanten-Hall-Systeme’’ and CFN of DFG, by US DOE under contract No. DE-AC02-98CH1-886, by the Program ‘‘Russian Science School’’ under grant No. 2192.2003.2, and by RFBR.

APPENDIX A: LUTTINGER FORMALISM FOR THE THERMODYNAMIC POTENTIAL

In this Appendix we derive the oscillatory part of the thermodynamic potential Ω , following Ref. 34. We calculate the sum in Eq. (7), using Poisson's summation formula

$$\sum_{m=0}^{\infty} f(m\omega_c) = \frac{1}{2}f(0) + \int_0^{\infty} \frac{d\xi}{\omega_c} f(\xi) - \sum_{k=1}^{\infty} \frac{1}{2\pi i k} \int_0^{\infty} d\xi f'(\xi) \left(e^{2\pi i k \xi / \omega_c} - e^{-2\pi i k \xi / \omega_c} \right). \quad (\text{A.1})$$

From Eq. (7) we have $f'(\xi) = -G(i\varepsilon_n, \xi)$, where $G(i\varepsilon_n, \xi)$ is the Green's function. Thus, to extract the oscillatory contribution to Ω , we should calculate the following integral

$$I_{\pm} = \int_0^{\infty} d\xi \frac{e^{\pm 2\pi i k \xi / \omega_c}}{\xi - \mu - i\varepsilon_n + \Sigma_{ee}(i\varepsilon_n, \xi) - i \operatorname{sgn} \varepsilon_n / 2\tau} \left[1 + \frac{\partial \Sigma_{ee}(i\varepsilon_n, \xi)}{\partial \xi} \right]. \quad (\text{A.2})$$

Here we introduce the self-energy $\Sigma_{ee}(i\varepsilon_n, \xi)$, which is a function defined in the plane of a complex variable ξ , such that

$$\Sigma_{ee}(i\varepsilon_n, m\omega_c) = \Sigma_{ee}(i\varepsilon_n, \xi = m\omega_c). \quad (\text{A.3})$$

The main contribution comes from the pole $\xi = \xi_0$, where ξ_0 obeys the self-consistent equation

$$\xi_0 = \mu + i\varepsilon_n + \frac{i}{2\tau} \operatorname{sgn} \varepsilon_n - \Sigma_{ee}(i\varepsilon_n, \xi_0). \quad (\text{A.4})$$

We expand the self-energy (from now on we will skip the subscript “ee” in Σ_{ee}) in the vicinity of the pole,

$$\Sigma(i\varepsilon_n, \xi) \simeq \Sigma(i\varepsilon_n, \xi_0) + (\xi - \xi_0) \frac{\partial \Sigma(i\varepsilon_n, \xi)}{\partial \xi} \Big|_{\xi=\xi_0}. \quad (\text{A.5})$$

Then the denominator in Eq. (A.2) becomes proportional to $(1 + \partial \Sigma / \partial \xi)$ so that these factors drop out and the integral takes a simple form

$$I_{\pm} = \int_0^{\infty} d\xi \frac{\exp[\pm 2\pi i k \xi / \omega_c]}{\xi - \xi_0}. \quad (\text{A.6})$$

We first single out the FL renormalization factors in Σ , i.e. represent the self-energy in the following form (assuming a constant electron concentration)

$$\Sigma(i\varepsilon_n, \xi) \simeq \delta\mu + \beta_0(\xi - \tilde{\mu}) - i\alpha_0\varepsilon_n + \delta\Sigma(i\varepsilon_n, \xi), \quad (\text{A.7})$$

where

$$\tilde{\mu} = \mu - \delta\mu = \pi n_e / m$$

is the chemical potential for noninteracting electrons (n_e is the electron concentration): $\xi - \tilde{\mu} \simeq v_F(k - k_F)$. The correction to the self-energy $\delta\Sigma(i\varepsilon_n, \xi)$ contains contributions that are smaller than $\beta_0(\xi - \mu)$ and $\alpha_0\varepsilon_n$ by either T/E_F or $1/E_F\tau$. These additional contributions are related to the inelastic processes and to the modification of the pure FL result due to disorder.

In order to solve Eq. (A.4), we treat the subleading terms constituting $\delta\Sigma$ as small corrections. Solving Eq. (A.4) by iterations, we first find its solution neglecting these small corrections and expanding Σ_{00} around the mass-shell:

$$\xi_0^{(0)} \simeq \tilde{\mu} + i\varepsilon_n + \frac{i}{2\tau} \operatorname{sgn} \varepsilon_n + i\alpha_0\varepsilon_n - \beta_0(\xi_0^{(0)} - \tilde{\mu}), \quad (\text{A.8})$$

which yields

$$\xi_0^{(0)} = \frac{\pi n_e}{m} + i\varepsilon_n \frac{1 + \alpha_0}{1 + \beta_0} + \frac{i \operatorname{sgn} \varepsilon_n}{2\tau(1 + \beta_0)}. \quad (\text{A.9})$$

Next we use this value of ξ_0 in $\delta\Sigma$ and solve the self-consistent equation again, now keeping the terms previously neglected. Then we arrive at

$$\xi_0 \simeq \xi_0^{(0)} - \delta\Sigma(i\varepsilon_n, \xi_0^{(0)}) / (1 + \beta_0). \quad (\text{A.10})$$

Substituting this value of the pole in Eq. (A.6), we obtain

$$\Omega = \Omega_{\text{osc}} + \Omega_{\text{Non-osc}}, \quad (\text{A.11})$$

where

$$\begin{aligned} \Omega_{\text{Non-osc}} = & -T\nu\omega_c \sum_{n=-\infty}^{\infty} \ln \left(-\mu - i\varepsilon_n + \Sigma(i\varepsilon_n, 0) - \frac{i \operatorname{sgn} \varepsilon_n}{2\tau} \right) \\ & - 2\nu T \sum_{n=-\infty}^{\infty} \int_0^{\infty} d\xi \ln \left(\xi - \mu - i\varepsilon_n + \Sigma(i\varepsilon_n, \xi) - \frac{i \operatorname{sgn} \varepsilon_n}{2\tau} \right) \end{aligned} \quad (\text{A.12})$$

and

$$\begin{aligned} \Omega_{\text{osc}} = & \sum_{k=1}^{\infty} \sum_{n=-\infty}^{\infty} \frac{T 2\nu\omega_c}{k} \exp \left\{ \frac{2\pi k}{\omega_c} \left(i \frac{\pi n_e}{m} \operatorname{sgn} \varepsilon_n - \frac{|\varepsilon_n|(1+\alpha_0) + 1/2\tau + i\delta\Sigma(i\varepsilon_n, \xi_0) \operatorname{sgn} \varepsilon_n}{1+\beta_0} \right) \right\} \\ = & \sum_{k=1}^{\infty} \frac{4\nu T\omega_c}{k} \cos \frac{2\pi^2 k n_e}{eB} \sum_{n=0}^{\infty} \exp \left\{ -\frac{2\pi k}{\omega_c} \frac{1+\alpha_0}{1+\beta_0} \left[\varepsilon_n + \frac{1}{2\tau(1+\alpha_0)} + \frac{i\delta\Sigma(i\varepsilon_n, \xi_0)}{1+\alpha_0} \right] \right\} \end{aligned} \quad (\text{A.13})$$

$$\simeq 2\nu \left(\frac{\omega_c}{2\pi} \right)^2 A_1 \cos \frac{2\pi^2 n_e}{eB}. \quad (\text{A.14})$$

Here A_1 is the amplitude of the principal harmonics of the oscillations,

$$A_1 \equiv \frac{4\pi^2 T}{\omega_c} \sum_{\varepsilon_n > 0} \exp \left(-\frac{2\pi}{\omega_c} \frac{1+\alpha_0}{1+\beta_0} \left[\varepsilon_n + \frac{1}{2\tau(1+\alpha_0)} + \frac{i\delta\Sigma(i\varepsilon_n, \xi_0)}{1+\alpha_0} \right] \right). \quad (\text{A.15})$$

APPENDIX B: CALCULATION OF SELF-ENERGIES

In this Appendix we calculate the relevant self-energy contributions and derive Eq. (28). The zero- B Green's function is given by

$$\begin{aligned} G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q}) &= \frac{i}{(\omega_m - \varepsilon_n)(1+\alpha_0) + \operatorname{sgn}(\omega_m - \varepsilon_n)/2\tau - [i(\xi - \tilde{\mu}) - iv_F q \cos \theta](1+\beta_0)} \\ &= \frac{i}{\tilde{\omega}_m - \tilde{\varepsilon}_n + \operatorname{sgn}(\omega_m - \varepsilon_n)/2\tau - i(\xi - \tilde{\mu})(1+\beta_0) + iv_F \tilde{q} \cos \theta}. \end{aligned} \quad (\text{B.1})$$

We denote the FL-renormalized energies and momenta as $\tilde{\omega}_m = (1+\alpha_0)\omega_m$, $\tilde{\varepsilon}_n = (1+\alpha_0)\varepsilon_n$, and $\tilde{q} = (1+\beta_0)q$. As the first approximation, we have set in Eq. (B.1) $p/m = v_F$ in the linear-in- q term $iv_F q \cos \theta$ and neglected $q^2/2m$.

Since the effective interaction (71) and (73) does not depend on the polar angle of the transferred momentum $\mathbf{q}$, we average the FL-dressed Green's function $G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q})$ over the angle between $\mathbf{p}$ and $\mathbf{q}$. The result of angle-averaging is

$$\langle G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q}) \rangle = \int \frac{d\theta}{2\pi} G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q}) \quad (\text{B.2})$$

$$= \frac{i \operatorname{sgn}(\omega_m - \varepsilon_n)}{\sqrt{[\tilde{\omega}_m - \tilde{\varepsilon}_n + \operatorname{sgn}(\omega_m - \varepsilon_n)/2\tau - i(\xi - \tilde{\mu})(1+\beta_0)]^2 + v_F^2 \tilde{q}^2}}. \quad (\text{B.3})$$

We substitute $\xi = \xi_0 \simeq \xi_0^{(0)} = \tilde{\mu} + i\tilde{\varepsilon}_n/(1+\beta_0) + i \operatorname{sgn} \varepsilon_n/2\tau(1+\beta_0)$ for ξ in (B.3) since we are interested in $\delta\Sigma(i\varepsilon_n, \xi_0)$ (the only place where ξ_0 appears in $\delta\Sigma$ is the Green's function under the interaction line). Then the denominator in (B.3) for $q = 0$ reads

$$\begin{aligned} & \tilde{\omega}_m - \tilde{\varepsilon}_n + \operatorname{sgn}(\omega_m - \varepsilon_n)/2\tau - i(\xi - \tilde{\mu})(1+\beta_0) \\ &= \tilde{\omega}_m - \tilde{\varepsilon}_n + \operatorname{sgn}(\omega_m - \varepsilon_n)/2\tau - i(1+\beta_0) \left[i \frac{\tilde{\varepsilon}_n}{1+\beta_0} + i \frac{\operatorname{sgn} \varepsilon_n}{2\tau(1+\beta_0)} \right] \\ &= \tilde{\omega}_m + \frac{1}{\tau} \theta(\omega_m - \varepsilon_n) \theta(\varepsilon_n), \end{aligned} \quad (\text{B.4})$$

with $\theta(x)$ the theta-function. For definiteness, below we consider $\varepsilon_n > 0$.

For Σ_{01} and Σ_{11} at $\varepsilon_n > 0$ we consider the Green's function at $\omega_m - \varepsilon_n > 0$ in order to have different signs of Matsubara energies in the Green's functions connected by the interaction vertex. This condition allows us to dress the interaction vertices by impurity ladders.

We see that ε_n drops out in the averaged Green's function taken at $\xi = \xi_0$, as in two-particle quantities:

$$\langle G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q}) \rangle |_{\xi=\xi_0} = \frac{i}{\sqrt{(|\tilde{\omega}_m| + 1/\tau)^2 + v_F^2 \tilde{q}^2}}, \quad \omega_m > \varepsilon_n. \quad (\text{B.5})$$

When both ε_n and $\varepsilon_n - \omega_m$ have the same sign (such a contribution appears in the calculation of $\delta\Sigma_{00}$), we find

$$\langle G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q}) \rangle |_{\xi=\xi_0} = -\frac{i}{\sqrt{|\tilde{\omega}_m|^2 + v_F^2 \tilde{q}^2}}, \quad \omega_m < \varepsilon_n. \quad (\text{B.6})$$

Now we re-define the Fermi velocity to absorb the FL-factors according to

$$v_F^* = v_F \frac{1 + \beta_0}{1 + \alpha_0} = \frac{k_F}{m^*}. \quad (\text{B.7})$$

Then we can return from $\tilde{\omega}_m$ and $\tilde{q}$ to ω_m and q , expressing the angle-averaged Green's function in terms of FL-renormalized parameters Z , τ^* [introduced in Eq. (15)] and v_F^* :

$$\begin{aligned} \langle G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q}) \rangle |_{\xi=\xi_0} &= \frac{iZ}{\sqrt{(|\omega_m| + 1/\tau^*)^2 + (v_F^* q)^2}}, \quad \omega_m > \varepsilon_n, \\ \langle G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q}) \rangle |_{\xi=\xi_0} &= -\frac{iZ}{\sqrt{|\omega_m|^2 + (v_F^* q)^2}}, \quad \omega_m < \varepsilon_n. \end{aligned}$$

Furthermore, the Z -factor will be cancelled in the final result, when $\langle G(i\varepsilon_n - i\omega_m, \vec{p} - \vec{q}) \rangle$ is used to calculate the correction to the observables, see e.g. Ref. 39 and discussion in Sec. IIB.

Using (B.5) and (B.6), we obtain

$$\begin{aligned} \Sigma_{01}^a(i\varepsilon_n, \xi_0) &= -T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} V(i\omega_m, \mathbf{q}) \Gamma(i\omega_m, \mathbf{q}) \langle G(i\varepsilon_n, \mathbf{p} - \mathbf{q}) \rangle |_{\xi_p=\xi_0} \\ &= -i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} \frac{V(i\omega_m, \mathbf{q}) \Gamma(i\omega_m, \mathbf{q})}{S(i\omega_m, \mathbf{q})}, \end{aligned} \quad (\text{B.8})$$

$$\begin{aligned} \Sigma_{11}^a(i\varepsilon_n, \xi_0) &= -T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} V(i\omega_m, \mathbf{q}) \Gamma^2(i\omega_m, \mathbf{q}) \langle G(i\varepsilon_n, \mathbf{p} - \mathbf{q}) \rangle |_{\xi_p=\xi_0} \\ &= -i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} \frac{V(i\omega_m, \mathbf{q}) \Gamma^2(i\omega_m, \mathbf{q})}{S(i\omega_m, \mathbf{q})}, \end{aligned} \quad (\text{B.9})$$

where for brevity we introduce new variables $S(i\omega_m, \mathbf{q}) \equiv \sqrt{(|\omega_m| + 1/\tau)^2 + v_F^2 q^2} = \sqrt{W^2 + v_F^2 q^2}$ and $W \equiv |\omega_m| + 1/\tau$.

Now we consider the contribution to the self-energy without vertex corrections, $\Sigma_{00}^a(i\varepsilon_n, \xi)$. We recall that in $\Sigma_{00}^a(i\varepsilon_n, \xi)$ the summation over transferred frequencies is not restricted to $\omega_m > \varepsilon_n$. Presenting $\Sigma_{00}^a(i\varepsilon_n, \xi)$ as

$$\Sigma_{00}^a(i\varepsilon_n, \xi_0) = \Sigma_{00}^{a,+,-}(i\varepsilon_n, \xi_0) + \Sigma_{00}^{a,++}(i\varepsilon_n, \xi_0), \quad (\text{B.10})$$

$$\begin{aligned} \Sigma_{00}^{a,+,-}(i\varepsilon_n, \xi_0) &= -T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} V(i\omega_m, \mathbf{q}) \langle G(i\varepsilon_n, \mathbf{p} - \mathbf{q}) \rangle |_{\xi_p=\xi_0} \\ &= -i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} \frac{V(i\omega_m, \mathbf{q})}{S(i\omega_m, \mathbf{q})}, \end{aligned} \quad (\text{B.11})$$

we further split the contribution $\Sigma_{00}^{a,++}(i\varepsilon_n, \xi_0)$ corresponding to no change of Matsubara frequencies at the interaction vertices into two parts as follows

$$\begin{aligned} \Sigma_{00}^{a,++}(i\varepsilon_n, \xi_0) &= -T \sum_{\omega_m < \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} V(i\omega_m, \mathbf{q}) \langle G(i\varepsilon_n, \mathbf{p} - \mathbf{q}) \rangle |_{\xi_p=\xi_0} \\ &= i T \int \frac{d^2 q}{(2\pi)^2} \left\{ \sum_{|\omega_m| < \varepsilon_n} + \sum_{\omega_m < -\varepsilon_n} \right\} \frac{V(i\omega_m, \mathbf{q})}{\sqrt{|\omega_m|^2 + q^2 v_F^2}} \end{aligned} \quad (\text{B.12})$$

The terms with $|\omega_m| < \varepsilon_n$ are responsible for the FL-renormalization and for the inelastic (determined by real processes) contribution to the self-energy, yielding the following FL-type term²⁶:

$$\Sigma^{\text{FL}}(i\varepsilon_n, \xi_0) = -i\alpha_0 \varepsilon_n - i\gamma(i\varepsilon_n, T) \frac{\varepsilon_n^2 - \pi^2 T^2}{E_F}, \quad (\text{B.13})$$

where the function $\gamma(i\varepsilon_n, T)$ depends logarithmically on $\max[\varepsilon_n, T, 1/\tau]$: in particular, $\gamma(i\varepsilon_n, T) \propto \ln[E_F/(-i\varepsilon_n)]$ for $\varepsilon_n \gg T, 1/\tau$ and $\gamma(i\varepsilon_n, T) \propto \ln[iE_F\tau]$ for $1/\tau \gg T, \varepsilon_n$. The first term in (B.13) determines the FL Z -factor and has been separated from $\delta\Sigma$ which governs the correction to the damping factor, see Eq. (A.7). As for the second term, its imaginary part describes the inelastic electron-electron scattering, while its real part contributes to the renormalization of the effective mass^{32,33,45}. However, when taken at $\varepsilon_0 = \pi T$, as appropriate for the damping of the magnetooscillations at $T \gg \omega_c$, the second term in (B.13) vanishes, in agreement with Ref. 26. Note that for the case of weak short-range interaction, $V(i\omega_m, \mathbf{q}) = \text{const}(\omega_m)$, the inelastic contribution is zero to the first order in V .

Thus in order to calculate $\delta\Sigma$, we shall retain in (B.12) only the term corresponding to the summation over $\omega_m < -\varepsilon_n$. In this term we change the sign of ω_m and (suppressing the irrelevant inelastic term) and obtain

$$\Sigma_{00}^{a,++}(i\varepsilon_n, \xi_0) = i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} \frac{V(i\omega_m, \mathbf{q})}{\sqrt{|\omega_m|^2 + q^2 v_F^2}}, \quad (\text{B.14})$$

thus arriving at

$$\Sigma_{00}^a(i\varepsilon_n, \xi_0) = -i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} V(i\omega_m, \mathbf{q}) \left[\frac{1}{S(i\omega_m, \mathbf{q})} - \frac{1}{S_0(i\omega_m, \mathbf{q})} \right]. \quad (\text{B.15})$$

Here we use the fact that $V(i\omega_m, \mathbf{q}) = V(-i\omega_m, \mathbf{q})$ as the dynamically screened interactions depends on ω_m only through $|\omega_m|$ and introduced $S_0(i\omega_m, \mathbf{q}) = \sqrt{\omega_m^2 + q^2 v_F^2}$.

Let us turn now to the Hikami-box diagrams, shown in Fig. 2b. We remind the reader that these diagrams are generated by covering each contribution to the self-energy Σ_{ij} from Fig. 2a with $\varepsilon_n(\varepsilon_n - \omega_m) < 0$ by a single impurity line. Therefore for white-noise disorder (addressed in this paper) the Hikami-box contribution to the self-energy is independent of ξ_p and can be expressed through the corresponding Σ_{ij}^a as

$$\Sigma_{ij}^b(i\varepsilon_n) = \frac{1}{2\pi\nu\tau} \int d\xi_p \nu [G(i\varepsilon_n, \xi_p)]^2 \Sigma_{ij}^a(i\varepsilon_n, \xi_p) \quad (\text{B.16})$$

$$= \frac{i}{\tau} \frac{\partial}{\partial \xi_p} \Sigma_{ij}^a(i\varepsilon_n, \xi_p) \Big|_{\xi_p = \xi_0} \quad (\text{B.17})$$

$$\Sigma_{ij}^b(i\varepsilon_n) = \frac{i}{\tau} \frac{\partial}{\partial \xi_p} \Sigma_{ij}^a(i\varepsilon_n, \xi_p) \Big|_{\xi_p = \xi_0} \quad (\text{B.18})$$

since the pole of G is given by ξ_0 , see Eq. (A.10). Differentiating (B.3) we get

$$\left\langle \frac{\partial}{\partial \xi_p} G(i\varepsilon_n - i\omega_n, \mathbf{p} - \mathbf{q}) \right\rangle \Big|_{\xi_p = \xi_0} = - \frac{|\omega_m| + 1/\tau}{\{(|\omega_m| + 1/\tau)^2 + v_F^2 q^2\}^{3/2}} = - \frac{W}{S^3}. \quad (\text{B.19})$$

This yields

$$\Sigma_{00}^b(i\varepsilon_n, \xi_0) = i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} \frac{V(i\omega_m, \mathbf{q}) W}{\tau S^3(i\omega_m, \mathbf{q})} \quad (\text{B.20})$$

$$\Sigma_{01}^b(i\varepsilon_n, \xi_0) = i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} \frac{V(i\omega_m, \mathbf{q}) \Gamma(i\omega_m, \mathbf{q}) W}{\tau S^3(i\omega_m, \mathbf{q})} \quad (\text{B.21})$$

$$\Sigma_{11}^b(i\varepsilon_n, \xi_0) = i T \sum_{\omega_m > \varepsilon_n} \int \frac{d^2 q}{(2\pi)^2} \frac{V(i\omega_m, \mathbf{q}) \Gamma^2(i\omega_m, \mathbf{q}) W}{\tau S^3(i\omega_m, \mathbf{q})} \quad (\text{B.22})$$

Combining all the contributions together, we arrive at Eq. (28).

APPENDIX C: COULOMB INTERACTION: TRIPLET CHANNEL

In this Appendix we calculate the contribution of the triplet channel to the damping of magnetooscillations. The effective interaction in the triplet channel can be found by replacing $V_0(q) \rightarrow F_0^\sigma/2\nu$ in the expression for the singlet channel:

$$\begin{aligned} 2\nu V^\sigma(i\omega_m, q) &= \frac{F_0^\sigma}{1 + F_0^\sigma \Pi(i\omega_m, q)} = \frac{F_0^\sigma}{1 + F_0^\sigma [1 - |\omega_m|/(S - 1/\tau)]} \\ &= \frac{F_0^\sigma (S - 1/\tau)}{(1 + F_0^\sigma)(S - 1/\tau) - |\omega_m| F_0^\sigma} = \frac{F_0^\sigma}{1 + F_0^\sigma} \frac{S - 1/\tau}{S - w}, \end{aligned} \quad (\text{C.1})$$

where we introduced

$$w \equiv |\omega_m| \frac{F_0^\sigma}{1 + F_0^\sigma} + \frac{1}{\tau}. \quad (\text{C.2})$$

This yields

$$2\nu V^\sigma(i\omega_m, q) K(i\omega_m, q) = \frac{F_0^\sigma}{1 + F_0^\sigma} \left[\frac{S_0 - (S - 1/\tau)}{S_0(S - w)} + \frac{S - W}{S - w} \frac{1}{S\tau(S - 1/\tau)} \right]. \quad (\text{C.3})$$

Performing the integration over q in (27) and taking into account the three triplet terms corresponding to different projection of the total spin on the z -axis $S_z = 0, \pm 1$, we obtain

$$\delta\Sigma^\sigma(i\pi T, \xi_0) = -\frac{i T}{2E_F\tau} \frac{3F_0^\sigma}{1 + F_0^\sigma} \sum_{\omega_m=2\pi T}^{\Delta} \left\{ \left(\frac{\omega_m}{\sqrt{\Delta^2 + \omega_m^2}} - 1 \right) + \ln \frac{\Delta^2 + \omega_m^2}{\omega_m^2} + h(\omega_m\tau, F_0^\sigma) \right\}, \quad (\text{C.4})$$

$$h(z, y) \simeq \left(1 - \frac{1}{y} \right) \ln[1 + y] + \frac{y}{1 + y} z h_1(x, y) \quad (\text{C.5})$$

$$\begin{aligned} h_1(z, y) &= \ln \frac{(1 + 2z)(1 + y)}{2z} + \frac{zy + (1 + y)}{\sqrt{|2(1 + y)z - (zy)^2|}} \\ &\times \begin{cases} \left[\arcsin \frac{zy - 1}{\sqrt{2z + 1}} - \arcsin \frac{zy + (1 + y)}{(1 + y)\sqrt{2z + 1}} \right], & z < \frac{2(1 + y)}{y^2}, \\ \left[\ln \left(1 + \frac{2 + y}{zy - 1 + \sqrt{(zy)^2 - 2(1 + y)z}} \right) - \ln(1 + y) \right], & z > \frac{2(1 + y)}{y^2}. \end{cases} \end{aligned} \quad (\text{C.6})$$

We see that the first two terms in (C.4) correspond to the point-like interaction with $\nu U_0 \rightarrow 3F_0^\sigma/(1 + F_0^\sigma)$, see Eq. (39). The term $h(\omega_m, F_0^\sigma)$ corresponds to the crossover function $f(x)$ in the singlet channel, see Eqs. (81) and (82). The result for the singlet channel is reproduced in the limit $F_0^\sigma \rightarrow \infty$ (cf. Ref. 9).

The summation over Matsubara frequencies leads to

$$\delta\Sigma^\sigma(i\pi T, \xi_0) = -\frac{i T}{2E_F\tau} \frac{3F_0^\sigma}{1 + F_0^\sigma} \left\{ \frac{\text{const}\Delta}{T} - \left[1 - \frac{\lambda(F_0^\sigma)}{8\pi T\tau} \right] \ln \frac{\Delta}{T} - f_\sigma(4\pi T\tau, F_0^\sigma) \right\}, \quad (\text{C.7})$$

where

$$\lambda(y) = \frac{1 + y}{y^3} [y(6 + y) - 2(3 + 2y) \ln(1 + y)] \quad (\text{C.8})$$

and

$$f_\sigma(x, y) = \sum_{m=1}^{\infty} \left\{ -h\left(\frac{mx}{2}, y\right) + \left[1 - \frac{2 \ln(1 + y)}{y} \right] + \frac{\lambda(y)}{2mx} \right\} \quad (\text{C.9})$$

As a result we obtain the following T dependence of the triplet contribution to the damping exponent

$$B^\sigma(T) = \frac{\pi}{\omega_c \tau} \frac{3F_0^\sigma}{1 + F_0^\sigma} \frac{T}{E_F} \left\{ \left[1 - \frac{\lambda(F_0^\sigma)}{8\pi T \tau} \right] \ln \frac{\Delta}{T} + f_\sigma(4\pi T \tau, F_0^\sigma) \right\}. \quad (\text{C.10})$$

For not too strong interaction, $(F_0^\sigma)^2/(1 + F_0^\sigma) \lesssim 1$, (i.e. for $|F_0^\sigma| \lesssim 0.6$, which is typically met in experiments, see e.g. Ref. 48) the crossover function $f_\sigma(x, y)$ only yields the subleading T -dependence⁴⁹ of $B^\sigma(T)$, so that the leading contribution to the damping is given by Eq. (42) for the short-range interaction with νU_0 replaced by $3F_0^\sigma/(1 + F_0^\sigma)$.

-
- [†] Also at A.F. Ioffe Physical-Technical Institute, 194021 St. Petersburg, Russia.
- [#] Also at Petersburg Nuclear Physics Institute, 188350 St. Petersburg, Russia.
- ¹ B.L. Altshuler and A.G. Aronov, in *Electron-electron interactions in disordered conductors*, edited by A.L. Efros and M. Pollak (Elsevier, 1985), p. 1.
- ² for a review see A.M. Finkelstein, Sov. Sci. Rev. A. Phys **14**, 1 (1990).
- ³ S.V. Kravchenko, G.V. Kravchenko, J.E. Furneaux, V.M. Pudalov, M. D'Iorio, Phys. Rev. B **50**, 8039 (1994); S.V. Kravchenko, W.E. Mason, G.E. Bowker, J.E. Furneaux, V.M. Pudalov, M. D'Iorio, Phys. Rev. B **51**, 7038 (1995); S.V. Kravchenko, D. Simonian, M.P. Sarachik, W. Mason, and J.E. Furneaux, Phys. Rev. Lett. **77**, 4938 (1996).
- ⁴ E. Abrahams, S.V. Kravchenko, and M.P. Sarachik, Rev. Mod. Phys. **73**, 251 (2001).
- ⁵ B.L. Altshuler, D.L. Maslov, and V.M. Pudalov, Physics E **9**, 209 (2001).
- ⁶ S.V. Kravchenko, and M.P. Sarachik, Rep. Prog. Phys. **67**, 1 (2004).
- ⁷ V.M. Pudalov, cond-mat/0405315.
- ⁸ F. Stern, Phys. Rev. Lett. **44**, 1469 (1980); A. Gold and V.T. Dolgoplov, Phys. Rev. B **33**, 1076 (1986); S. Das Sarma, Phys. Rev. B **33**, 5401 (1986); S. Das Sarma and E.H. Hwang, Phys. Rev. Lett. **83**, 164 (1999).
- ⁹ G. Zala, B.N. Narozhny, and I.L. Aleiner, Phys. Rev. B **64**, 214204 (2001); Phys. Rev. B **64**, 201201 (2001); Phys. Rev. B **65**, 020201 (2002).
- ¹⁰ I.V. Gornyi and A.D. Mirlin, Phys. Rev. Lett. **90**, 076801 (2003); Phys. Rev. B **69**, 045313 (2004).
- ¹¹ A. Punnoose and A.M. Finkelstein, Phys. Rev. Lett. **88**, 016802 (2002); Science **310**, 289 (2005).
- ¹² T. Okamoto, K. Hosoya, S. Kawaji, and A. Yagi, Phys. Rev. Lett. **82**, 3875 (1999).
- ¹³ S.V. Kravchenko, A.A. Shashkin, D.A. Bloore, and T.M. Klapwijk, Solid State Commun. **116**, 495 (2000).
- ¹⁴ V.M. Pudalov, M.E. Gershenson, H. Kojima, N. Butch, E.M. Dizhur, G. Brunthaler, A. Prinz, and G. Bauer, Phys. Rev. Lett. **88**, 196404 (2002).
- ¹⁵ J. Zhu, H.L. Stormer, L.N. Pfeiffer, K.W. Baldwin, and K.W. West, Phys. Rev. Lett. **90**, 056805 (2003).
- ¹⁶ S.A. Vitkalov, H. Zeng, K.M. Mertes, M.P. Sarachik, and T.M. Klapwijk, Phys. Rev. Lett. **87**, 086401 (2001).
- ¹⁷ A.A. Shashkin, S.V. Kravchenko, V.T. Dolgoplov, and T.M. Klapwijk, Phys. Rev. Lett. **87**, 086801 (2001).
- ¹⁸ O. Prus, Y. Yaish, M. Reznikov, U. Sivan, and V. Pudalov, Phys. Rev. B **67**, 205407 (2003).
- ¹⁹ V.M. Pudalov, M.E. Gershenson, and H. Kojima, in: *Fundamental Problems of Mesoscopic Physics. Interaction and Decoherence*, edited by I.V. Lerner, B.L. Altshuler, and Y. Gefen (Kluwer, 2004), p.309.
- ²⁰ A.A. Shashkin, Physics-Uspekhi **48**, 129 (2005).
- ²¹ A.A. Shashkin, S.V. Kravchenko, V.T. Dolgoplov, and T.M. Klapwijk, Phys. Rev. B **66**, 073303 (2002).
- ²² Y.Y. Proskuryakov, A.K. Savchenko, S.S. Safonov, M. Pepper, M.Y. Simmons, and D.A. Ritchie, Phys. Rev. Lett. **89**, 076406 (2002); Y.Y. Proskuryakov, A.K. Savchenko, S.S. Safonov, M. Pepper, M.Y. Simmons, D.A. Ritchie, E.H. Linfield, and Z.D. Kvon, J. Phys. A **36**, 9249 (2003).
- ²³ V.M. Pudalov, M.E. Gershenson, H. Kojima, G. Brunthaler, A. Prinz, and G. Bauer, Phys. Rev. Lett. **91**, 126403 (2003).
- ²⁴ S. Anissimova, A. Venkatesan, A.A. Shashkin, M.R. Sakr, S.V. Kravchenko, and T.M. Klapwijk, cond-mat/0503123.
- ²⁵ A.A. Shashkin, M. Rahimi, S. Anissimova, S.V. Kravchenko, V.T. Dolgoplov, and T.M. Klapwijk, Phys. Rev. Lett. **91**, 046403 (2003); A.A. Shashkin, S.V. Kravchenko, V.T. Dolgoplov, and T.M. Klapwijk, J. Phys. A: Math. Gen. **36**, 9237 (2003).
- ²⁶ G.M. Martin, D.L. Maslov, and M.Yu. Reizer, Phys. Rev. B **68**, 241309(R) (2003).
- ²⁷ I.M. Lifshitz and A.M. Kosevich, Zh. Eksp. Teor. Fiz. **29**, 730 (1955) [Sov. Phys. JETP **2**, 636 (1956)].
- ²⁸ S. Curnoe and P. C. E. Stamp, Phys. Rev. Lett. **80**, 3312, (1998).
- ²⁹ M. Fowler and R. E. Prange, Physics **1**, 315 (1965); S. Engelsberg and G. Simpson, Phys. Rev. B **2**, 1657 (1970).
- ³⁰ J. M. Luttinger and J. C. Ward Phys. Rev. **118**, 1417 (1960).
- ³¹ A.A. Abrikosov, L.P. Gorkov, and I.E. Dzyaloshinski, *Methods of quantum field theory in statistical physics*, (Dover Publications, New York, 1963).
- ³² A.V. Chubukov and D.L. Maslov, Phys. Rev. B **68**, 155113 (2003); Phys. Rev. B **69**, 121102 (2004).
- ³³ A.V. Chubukov, D.L. Maslov, S. Gangadharaiiah, and L.I. Glazman, Phys. Rev. B **71**, 205112 (2005); Phys. Rev. Lett. **95**, 026402 (2005); S. Gangadharaiiah, D.L. Maslov, A.V. Chubukov, and L.I. Glazman, Phys. Rev. Lett. **94**, 156407 (2005).
- ³⁴ J.M. Luttinger, Phys. Rev. **121**, 1251 (1961).
- ³⁵ T. Ando, A.B. Fowler, and F. Stern, Rev. Mod. Phys. **54**, 437 (1982).

- ³⁶ In strong magnetic field and smooth disorder this is no longer true since the transport acquires adiabatic character due to classical memory effects, see e.g. A.D. Mirlin, D.G. Polyakov, and P. Wölfle, Phys. Rev. Lett. **80**, 2429 (1998)
- ³⁷ I.V. Gornyi, A.D. Mirlin, and F. von Oppen, Phys. Rev. B **70**, 245302 (2004).
- ³⁸ A. M. Rudin, I. L. Aleiner, and L. I. Glazman, Phys. Rev. B **55**, 9322 (1997).
- ³⁹ G. Zala, B.N. Narozhny, I.L. Aleiner, and V.I. Fal'ko, Phys. Rev. B **69**, 075306 (2004).
- ⁴⁰ A.D. Mirlin, E. Altshuler, and P. Wölfle, Annalen der Physik **5**, 281 (1996).
- ⁴¹ G. Catelani and I.L. Aleiner, JETP **100**, 331 (2005).
- ⁴² M.G. Vavilov and I.L. Aleiner, Phys. Rev. B **69**, 035303 (2004).
- ⁴³ B.L. Altshuler, A.G. Aronov, and D.E. Khmelnitsky, J. Phys. C **15**, 7367 (1982).
- ⁴⁴ T. Ludwig and A.D. Mirlin, Phys. Rev. B **69**, 193306 (2004).
- ⁴⁵ V.M. Galitski and S. Das Sarma, Phys. Rev. B **70**, 035111 (2004).
- ⁴⁶ This calculation is in fact very similar to the analysis of the momentum dependence of the RKKY interaction between the localized moments via the conduction electrons, see D.N. Aristov, Phys. Rev. B **55**, 8064 (1997).
- ⁴⁷ Technically, this difference between the Coulomb interaction case and the short-range interaction case can be traced back to the contribution of Σ_{00} represented by the first term in the r.h.s. of Eq. (78). For Coulomb interaction this term produces in the diffusive regime a $T \ln T$ contribution to the damping factor. In the ballistic regime, where the frequency dependence of screening becomes irrelevant (*cf.* the conductivity corrections, Refs. 9,10) and the Coulomb interaction reduces to a constant, $V^p(i\omega_m, \mathbf{q}) \simeq 1/2\nu$, this term yields a subleading contribution as in the case of the short-range interaction.
- ⁴⁸ E.A. Galaktionov, A.K. Savchenko, S.S. Safonov, Y.Y. Proskuryakov, L. Li, M. Pepper, M.Y. Simmons, D.A. Ritchie, E.H. Linfield, and Z.D. Kvon, in *Fundamental Problems of Mesoscopic Physics: Interactions and Decoherence*, edited by I.V. Lerner, B.L. Altshuler, and Y. Gefen (Kluwer Academic Publishers, Dordrecht, 2004), p. 349.
- ⁴⁹ For strong interaction $(F_0^\sigma)^2/(1 + F_0^\sigma) \gg 1$ (in particular, in the vicinity of the ferromagnetic instability $F_0^\sigma \rightarrow -1$) an additional regime, $(1 + F_0^\sigma)/(F_0^\sigma)^2 \ll 2\pi T\tau \ll 1$, appears in the diffusive range of T .